



63426/B

The Library of the
Wellcome Institute for
the History of Medicine

MEDICAL SOCIETY
OF
LONDON
DEPOSIT

Accession Number

Press Mark

WARNER, R

From The Author

U₁

PLANTÆ WOODFORDIENSES
A
CATALOGUE
OF THE
MORE PERFECT
PLANTS
GROWING SPONTANEOUSLY ABOUT
WOODFORD

IN THE COUNTY OF

S E X.



LONDON
Printed for the AUTHOR. 1771.

Κατανοήσατε τὰ κῆρα, πῶς αὐξάνει
οὐ κοπιᾷ οὐδὲ νήθει. λέγω δὲ ὑμῖν οὐδὲ
Σολομῶν ἐν πάσῃ τῇ δόξῃ αὐτοῦ πε-
ριεβάλετο ὡς ἐν τούτων.

S. Luc. xii. 27.

TO
MR. JOHN LISLE
MASTER
MR. JOHN CHANNING
AND
MR. JOHN PEARCE
WARDENS
AND THE REST OF
THE COURT OF ASSISTANTS
OF
THE WORSHIPFUL COMPANY
OF
APOTHECARIES OF LONDON
THIS CATALOGUE
IN A GRATEFUL SENSE OF THEIR FAVOUR;
IN THE PROSECUTION OF HIS
BOTANICAL AMUSEMENTS
IS
INSCRIBED
BY
THEIR OBLIGED HUMBLE SERVANT
RICHARD WARNER.



P R E F A C E.

WHAT is here offered to the lovers of Botany, is the result of several investigations for some years past, of the more perfect Plants growing spontaneously about WOODFORD, in the COUNTY of E S S E X, the residence of The **AUTHOR** almost from infancy. In the prosecution of which, he hereby gratefully acknowledges the assistance of the following Gentlemen, his Friends and Companions in his researches for that purpose :

Mr. *Stainsby Alchorne.*

Mr. *John Bradney.*

Mr. *Joseph Bradney.*

Mr. *John Channing.*

Mr. *John Channing junr.*

Mr. *John Carwte.*

Mr. *Josiah Colebrooke.*

Mr. *James Gordon.*

Mr. *William Gordon.*

Mr.

Mr. *James Gordon junr.*
Mr. *Alexander Gordon.*
Mr. *William Hudson.*
Mr. *Samuel Longbotham.*
Mr. *Isaac Mather.*
Mr. *Philip Miller.*
Mr. *Peter Henry Norris.*
Mr. *Samuel Parker.*
Dr. *William Watson.*
Dr. *William Watson junr.*
Dr. *John Wilkinson.*

To which, are to be added, the late Mr. *Joseph Bradney*, and the late Dr. *John Wilmer.*

In printing them, he has followed the example of Mr. *Ray*, who published a Catalogue of Plants growing about CAMBRIDGE; Mr. *Blackstone*, of those round HAREFIELD; Mr. *Dering*, of those round NORTHAMPTON; and many others. And, though he does not know that many scarce or very uncommon Plants are to be found in this Circuit, yet, as it has been often hinted by Botanical Writers, that if an account was taken of the Plants growing about particular places, it might be a means of promoting Botanical Knowledge, and assisting young Students
in

in that Science, the following mite is offered for that purpose.

He has given two *Latin* Names to each Plant; the first, taken from RAY's *Synopsis Stirpium Britannicarum*, the third edition, 1724; the other from LINNÆUS, as quoted in HUDSON's *Flora Anglica*, edition 1762. He has added the Class assigned by LINNÆUS; given the *English* name or names, usually from RAY; mentioned the Soil wherein each Plant is found; very often the particular place of Growth, and the time of Flowering.

In order to render it of more general use to the *English* Reader, he has endeavoured to make it, as much as possible, an *English* Catalogue; by specifying the soil, the place of Growth, and the time of Flowering, in the *English* language.

If what is here offered should be of any use, for the very laudable purpose above-mentioned, his end in making it public, is abundantly answered.

Woodford-Row,
July 1, 1771.

RICHARD WARNER.

P L A N T Æ

W O O D F O R D I E N S E S.

A.

ABSINTHIUM vulgare. *Raii Syn.* 188.

ARTEMISIA [*Absinthium*] foliis compositis,
multifidis, floribus sub-globosis pendulis;
receptaculo villoso. *Hudsoni Fl.* 311.

Classis Linnæi Syngenesia: Polygamia Superflua.
Common Wormwood.

*Found upon dry banks near the road side, in
plenty.*

It flowers in August.

ACER majus. *Raii Syn.* 470.

ACER [*Pseudo-platanus*] foliis quinquelobis inæqualiter ferratis, floribus racemosis. *Hudsoni Fl.* 378.

Classis Linnæi Polygamia Monoecia.

The greater Maple, or Sycamore tree.

Found in hedges, near houses.

It flowers in May or June.

ACER minus. *Raii Syn.* 470.

ACER [*campestre*] foliis lobatis obtusis emarginatis. *Hudsoni Fl.* 379.

Classis Linnæi Polygamia Monoecia.

The common Maple.

In hedges almost every where; and in the woody parts of the Forest: not uncommon.

It flowers in May or June.

ACORUS verus, five Calamus officinarum. *Raii Syn.* 437.

ACORUS [*Calamus*] *Hudsoni Fl.* 128.

Classis Linnæi Hexandria Monogynia.

The sweet-smelling Flag, or Calamus.

Found on the sides of some ponds on the hill near Mr. Warner's Piece of water, opposite Harts, his house at Woodford Row; and in the river Rhodon, near Woodford Bridge, opposite the eight mile stone; but in general, not very common.

It flowers or produces its Iuli in May.

WOODFORDIENSES. 3

ADIANTUM nigrum officinarum. *Raii Syn.*
126.

ASPLENIUM [*Adiantum nigrum*] frondibus subtripinnatis, foliolis alternis, pinnis lanceolatis inciso ferratis. *Hudsoni Fl.* 386.

Classis Linnæi Cryptogamia Filices.

Common black Maiden Hair.

In shady lanes on the banks of hollow ways, and on old walls. In these parts not very common; but found on the buttresses on the North and South sides of Loughton Church, in plenty.

AGRIFOLIUM. *Raii Syn.* 466.

ILEX [*Aquifolium*] foliis ovatis acutis spinosis.
Hudsoni Fl. 505.

Classis Linnæi Tetrandia Trigynia.

The Holly tree.

In hedges, and on the Forest, in great plenty.
It flowers in May or June.

AGRIMONIA. *Raii Syn.* 202.

AGRIMONIA [*Eupatoria*] foliis caulinis pinnatis, foliolis undique ferratis, omnibus minutis interstinctis, fructibus hispidis. *Hudsoni Fl.* 180.

Classis Linnæi Dodecandria Digynia.

Agrimony.

On the sides of fields, and under hedges, in great plenty.

It flowers in June.

ALCEA vulgaris. *Raii Syn.* 252.

MALVA [*Alcea*] caule erecto, foliis multipar-
titis scabriusculis. *Hudsoni Fl.* 268.

Classis Linnæi Monodelphia Polyandria.

Vervain Mallow.

In meadows : not very uncommon.

It flowers in August.

ALNUS. *Raii Syn.* 442.

BETULA [*Alnus*] pedunculis ramosis. *Hud-
soni Fl.* 506.

Classis Linnæi Monoecia Triandria.

The common Alder tree.

*In moist places, and on the sides of ditches,
and banks of rivers, not uncommon ; on the
banks of the river Rhodon, between Chigwell
and Loughton, in plenty.*

It flowers in July.

ALLIUM sylvestre. *Raii Syn.* 369.

ALLIUM [*vineale*] caule teretifolio bulbifero,
staminibus tricuspидatis. *Hudsoni Fl.* 121.

Classis Linnæi Hexandria Monogynia.

Crow-garlick.

*In fields and meadows, and sometimes among
corn. Found in a field of Mr. Warner's behind
his house, in plenty.*

It flowers in June.

WOODFORDIENSES. 5

ALSINE hirsuta Myosotis. *Raii Syn.* 349.

CERASTIUM [*vulgatum*] foliis ovatis, petalis calyci æqualibus, caulibus diffusis. *Hudsoni Fl.* 175.

Classis Linnæi Decandria Pentagynia.

Narrow-leaved Mouse-ear Chickweed.

In fields and meadows : common.

It flowers in May or June.

ALSINE hirsuta Myosotis latifolia præcocior. *Raii Syn.* 348.

CERASTIUM [*viscosum*] erectum, villoso-viscosum. *Hudsoni Fl.* 175.

Classis Linnæi Decandria Pentagynia.

The broader leaved Mouse-ear Chick-weed.

In fields on a dry soil : not uncommon.

It flowers in April or May.

ALSINE longifolia, uliginosis proveniens locis. *Raii Syn.* 347.

STELLARIA [*graminea*] foliis linearibus integerrimis, floribus paniculatis. *Hudsoni Fl.* 166.

Classis Linnæi Decandria Trigynia.

Long leaved Water Chickweed.

On boggy grounds, and by rivulets and ditches that carry water all the year. Not very uncommon. Found on the bog between Snarebrook and the Woodford Bridge road.

It flowers in July.

ALSINE major repens perennis. *Raii Syn.*
347.

CERASTIUM [*aquaticum*] foliis cordatis sessilibus floribus solitariis, fructibus pendulis.
Hudsoni Fl. 177.

Classis Linnæi Decandria Pentagynia.
Great Marsh Chickweed.

In moist marshy places, and on the banks of the river Rhodon: not uncommon.

It flowers in July.

ALSINE minor multicaulis. *Raii Syn.* 349.

ARENARIA [*serpyllifolia*] foliis subovatis acutis sessilibus, corollis calyce brevioribus. *Hudsoni Fl.* 167.

Classis Linnæi Decandria Trigynia.
The least Chickweed.

On roofs of old houses, old walls, and on dry sandy soils: very common.

It flowers in May or June.

ALSINE parva palustris tricoccus portulacæ aquaticæ similis. *Raii Syn.* 353.

MONTIA [*fontana*] *Hudsoni Fl.* 48.

Classis Linnæi Triandria Digynia.

Small Water Chickweed or Purflane; by some called Blinks.

Near springs and on moist, marshy ground: not uncommon.

It flowers in March.

WOODFORDIENSES. 7

ALSINE Plantaginis folio. *Raii Syn.* 349.

ARENARIA [*trinervia*] foliis ovatis acutis, petiolatis, nervosis. *Hudsoni Fl.* 167.

Classis Linnæi Decandria Trigynia.

Plantain leaved Chickweed.

In moist woods and shady places: not uncommon.

It flowers in May or June.

ALSINE spergula dicta major. *Raii Syn.* 351.

SPERGULA [*arvensis*] foliis verticillatis, floribus decandris. *Hudsoni Fl.* 177.

Classis Linnæi Decandria Pentagynia.

Spurrey.

In corn fields on a dry soil: common.

It flowers in August.

ALSINELLA foliis caryophylleis. *Raii Syn.* 344.

SAGINA [*erecta*] caule erecto subunifloro.

Hudsoni Fl. 64.

Classis Linnæi Tetrandria Tetragynia.

The least Stitchwort.

On gravelly soils on the Forest: Found near Mr. Warner's great Piece of Water, commonly called The Gravel Pit: not common.

It flowers in April or May.

ALSINELLA muscoso flore repens. *Raii Syn.* 345.

SAGINA [*procumbens*] ramis procumbentibus.

Hudsoni Fl. 63.

Classis Linnæi Tetrandria Tetragynia.

Pin Pearlwort. Chickweed Breakstone.

On Walls, and in sandy ground: very common.

It flowers in June.

ANACAMPSEROS, *vulgò*, Faba crassa. *Raii*
Syn. 269.

SEDUM [*Telephium*] foliis planiusculis ferratis,
 corymbo folioso, caule erecto. *Hudsoni Fl.*
 170.

Classis Linnæi Decandria Pentagynia.

Orpine or Live-long.

In shady places under hedges, and sometimes in fields. Found on the side of a field between Muncombe and Luxborough-House, Sir Edward Walpole's : not common.

It flowers in August.

ANAGALLIS flore phoeniceo. *Raii Syn.* 282.

ANAGALLIS [*arvensis*] foliis indivisis, caule
 procumbente. *Hudsoni Fl.* 73.

Classis Linnæi Pentandria Monogynia.

Male Pimpernel.

In fields and gardens ; also among corn : common.

It flowers from May to August.

ANEMONE nemorum alba. *Raii Syn.* 259.

ANEMONE [*nemorosa*] feminibus acutis, foliolis
 incisis, caule unifloro. *Hudsoni Fl.* 208.

Classis Linnæi Polyandria Polygynia.

Wood Anemony.

In woods, under hedges, and all over the woody parts of the Forest : common.

It flowers in April.

WOODFORDIENSES. 9

ANGELICA sylvestris. *Raii Syn.* 208.

ANGELICA [*sylvestris*] foliis æqualibus, ovato-lanceolatis ferratis. *Hudsoni Fl.* 103.

Classis Linnæi Pentandria Digynia.

Wild Angelica.

In moist places, under hedges and near rivers.

It flowers in June or July.

ANGELICA sylvestris minor, seu erratica. *Raii Syn.* 208.

ÆGOPodium [*Podagraria*] foliis caulinis summis ternatis. *Hudsoni Fl.* 111.

Classis Linnæi Pentandria Digynia.

Herb Gerard, Gout-weed, or Ash-weed.

Under hedges, and common in gardens. Found in Mr. Warner's Garden, at Woodford Row, growing spontaneously, in great plenty.

It flowers in May.

ANONIS spinosa flore purpureo. *Raii Syn.* 332.

ONONIS [*spinosa*] floribus racemosis solitariis, foliis ternatis, ramis spinosis. *Hudsoni Fl.* 273.

Classis Linnæi Diadelphia Decandria.

Rest-harrow. Cammock. Petty Whin.

In dry fields and on the forest: common. Found in plenty in the foot-way from Woodford Row to Woodford Church, under the garden-wall belonging to the House commonly called, The Naked Beauty, Mr. Duffin's, in great plenty.

It flowers in June, July, or August.

APARINE. *Raii Syn.* 225.

GALIUM [*Aparine*] foliis octonis lanceolatis carinatis scabris retrorsum aculeatis, geniculis villosis, fructu hispido. *Hudsoni Fl.* 57.

Classis Linnæi Tetandria Monogynia.

Cleavers, or Goose-grafs.

Under hedges, in corn fields, and in gardens, spontaneously: very common.

It flowers from May to August.

APIUM palustre et officinarum. *Raii Syn.* 214.

APIUM [*grave olens*] foliis caulinis cuneiformibus. *Hudsoni Fl.* 111.

Classis Linnæi Pentandria Digynia.

Smallage.

In marshy places, and large ditches. Found by the road side from the eight mile stone to Woodford Bridge, and on the banks of the river Rhodon: not very common.

It flowers in August.

ANTIRRHINUM angustifolium sylvestre. *Raii Syn.* * 283.

ANTIRRHINUM [*orentium*] corollis caudatis, calycibus corolla longioribus. *Hudsoni Fl.* 239.

Classis Linnæi Didynamia Angiospermia.

The least Snap-dragon, or Calf's Snout.

In fields, on a dry, sandy soil, and amongst corn.

It flowers in July or August.

ANTIR-

WOODFORDIENSES. 11

ANTIRRHINUM majus. In *Raii Syn.* non invenitur.

ANTIRRHINUM [*majus*] corollis ecaudatis, calycibus rotundatis. *Hudsoni Fl.* 239.

Classis Linnæi Didynamia Angiospermia.

The greater Snap-dragon.

On old walls, and in chalk-pits. Found on the wall before Mr. Raymond's House, near Chigwell Church; and on the North Wall of Mr. Warner's Garden, near his House, in plenty: not uncommon.

It flowers from June to September.

ARTEMISIA vulgaris. *Raii Syn.* 190.

ARTEMISIA [*vulgaris*] foliis pinnatifidis planis incisis: subtus tomentosis, racemis simplicibus, floribus ovatis, radio quinquefloro.

Hudsoni Fl. 312.

Classis Linnæi Syngenesia: Polygamia superflua.
Mugwort.

Under Hedges, and on the sides of fields: common.

It flowers in August.

ARUM- *Raii Syn.* 266.

ARUM [*maculatum*] acaule, foliis hastatis integerrimis, spadice clavato. *Hudsoni Fl.* 342.

Classis Linnæi Gynandria Polyandria.

Wake Robin. Cuckow Point. Lords and Ladies.

In shady places, and under hedges: common.

It flowers in May.

ASPERULA. *Raii Syn.* 224.

ASPERULA [*odorata*] foliis octonis lanceolatis,
floribus fasciculatis pedunculatis. *Hudsoni*
Fl. 55.

Classis Linnæi Tetrandria Monogynia.

Woodroof.

In shady parts of the Forest: not very common.

It flowers in May.

ASPLENIUM five CETERACH. *Raii Syn.* 118.

ASPLENIUM [*Ceterach*] frondibus pinnatifidis;
lobis alternis confluentibus. *Hudsoni Fl.* 385.

Classis Linnæi Cryptogamia Filices.

Spleenwort, or Miltwaist.

*Found on a wall in Dr. Wilkinfon's Orchard,
the side of the lane leading to The Windmill;
and on an old wall belonging to Mr. Warner:
very uncommon.*

ATRIPLEX sylvestris angustifolia. *Raii Syn.*
151.

ATRIPLEX [*patula*] caule herbaceo patulo,
foliis sub-deltoideo-lanceolatis, calycibus fe-
minum disco dentatis. *Hudsoni Fl.* 377.

Classis Linnæi Polygamia Monoecia.

Narrow leaved wild Orach.

*In gardens, on dunghills, and in waste places:
common.*

It flowers in August.

WOODFORDIENSES. 13

ATRIPLEX sylvestris folio hastato, seu deltoide.

Raii Syn. 151.

ATRIPLEX [*hastata*] caule herbaceo, valvulis foemineis magnis deltoidibus sinuatis. *Hudsoni Fl.* 377.

Classis Linnæi Polygamia Monoecia.

Wild Orach, with a spear-pointed leaf.

In gardens, on dunghills, and in waste places : common.

It flowers in August or September.

B.

BALLOTE. *Raii Syn.* 244.

BALLOTA [*nigra*] cordatis indivisis ferratis, calycibus acuminatis. *Hudsoni Fl.* 227.

Classis Linnæi Didynamia Gymnospermia.

Stinking Horehound.

In barren places and under hedges: very common.

It flowers in August.

BARBAREA foliis minoribus et frequentius sinuatis. *Raii Syn.* 297.

ERYSIMUM [*Barbarea*] foliis lyratis, extimo subrotundo. *Hudsoni Fl.* 251. β.

Classis Linnæi Tetradynamia filiquosa.

Winter Cresses, or Rocket, with a small sinuated leaf.

In ditches and moist places: not very common.

It flowers in May.

BELLADONNA. *Raii Syn.* 265.

ATROPA [*Belladonna*] caule herbaceo, foliis ovatis integris. *Hudsoni Fl.* 79.

Classis Linnæi Pentandria Monogynia.

Deadly Nightshade, or Dwale.

Under hedges, and on rubbish. Found in the upper end of the Walk round Mr. Warner's fields under Capt. Foulis's garden-wall: not very common.

It flowers in June or July.

WOODFORDIENSES. 15

BELLIS sylvestris minor. *Raii Syn.* 184.

BELLIS [*perennis*] scapo nudo. *Hudsoni Fl.*

320.

Classis Linnæi Syngenefia: Polygamia superflua.
Common Daisie.

In grass walks in gardens, on the turf of lawns, and in meadows: very common.

It flowers almost the whole year.

BERBERIS dumetorum. *Raii Syn.* 465.

BERBERIS [*vulgaris*] pedunculis racemosis.

Hudsoni Fl. 119.

Classis Linnæi Hexandria Monogynia.

The Berberry, or, Pipperidge Bush.

Found in a lane leading from Dr. Wilkinson's to The Windmill; and in a lane near the five mile stone, in the road over Lee Bridge, which leads to Marsh-street, Walthamstow, in great plenty.

It flowers in May.

BETONICA. *Raii Syn.* 238.

BETONICA [*officinalis*] spica interrupta, corollarum lacinia labii intermedia emarginata.

Hudsoni Fl. 220.

Classis Linnæi Didynamia Gymnospermia.

Wood Betony.

In woods and under hedges; found occasionally, but not common, with a white flower.

It flowers in July or August.

BETULA.

BETULA. *Raii Syn.* 443.

BETULA [*alba*] foliis ovatis acuminatis ferratis.

Hudsoni Fl. 354.

Classis Linnæi Monoecia Tetrandria.

The Birch Tree.

Found on the Forest between High Beech, and Golden Hill in the Parish of Loughton : in general, not very common.

It flowers in April.

BLITUM Atriplex sylvestris dictum. *Raii Syn.* 154.

CHENOPODIUM [*album*] foliis rhomboideo-tri-angulis erosis posticè integris: summis oblongis, racemis erectis. *Hudsoni Fl.* 91.

Classis Linnæi Pentandria Digynia.

Common wild Orache.

Found in uncultivated places, on dunghills, and among rubbish: common.

It flowers in August.

BLITUM Ficus folio. *Raii Syn.* 155.

CHENOPODIUM [*serotinum*] foliis deltoides, sinuato-dentatis rugosis glabris uniformibus, racemis terminalibus. *Hudsoni Fl.* 91.

Classis Linnæi Pentandria Digynia.

Late flowering or Fig-leaved Blite.

On dunghills and among rubbish: not uncommon.

It flowers from August to October.

WOODFORDIENSES. 17

BLITUM perenne, *Bonus Henricus* dictum. *Raii Syn.* 156.

CHENOPODIUM [*Bonus Henricus*] foliis triangulari-sagittatis integerrimis. *Hudsoni Fl.* 89.
Classis Linnæi Pentandria Digynia.

Common English Mercury, or All-good.

Found on rubbish, old walls, by the road side, and, sometimes, in fields : not uncommon.

It flowers in August.

BLITUM *Pes anserinus* dictum. *Raii Syn.* 154.

CHENOPODIUM [*murale*] foliis acutis dentatis, racemis ramosis nudis. *Hudsoni Fl.* 90.

Classis Linnæi Pentandria Digynia.

Common Goose-foot, or Sow-bane.

On cultivated land ; and often, on rubbish : common.

It flowers in August.

BLITUM *Pes anserinus* dictum acutiore folio.
Raii Syn. 154.

CHENOPODIUM [*rubrum*] foliis cordato-triangularibus, obtusiusculis dentatis, racemis erectis compositis, subfoliosis caule brevioribus. *Hudsoni Fl.* 90.

Classis Linnæi Pentandria Digynia.

Sharp leaved Goose-foot, or Sow-bane.

On dunghills and rubbish : common.

It flowers in August.

BLITUM rubrum minus. *Raii Syn.* 157.

AMARANTUS [*Blitum*] spicis triandris erectiusculis, foliis ovatis retusis. *Hudsoni Fl.* 356.

Classis Linnæi Monoecia Pentandria.

The least Blite.

Among rubbish, on dunghills, and in gardens : very common.

It flowers in August.

BORRAGO hortensis. *Raii Syn.* 228.

BORAGO [*officinalis*] foliis omnibus alternis, calycibus patentibus. *Hudsoni Fl.* 68.

Classis Linnæi Pentandria Monogynia.

Borage.

Found on old walls, and by the road side : not uncommon. It is sometimes cultivated in gardens.

It flowers in May.

BRYONIA alba. *Raii Syn.* 261.

BRYONIA [*alba*] foliis palmatis, utrinque calloso-scabris. *Hudsoni Fl.* 373.

Classis Linnæi Dioecia Syngenesia.

White Briony.

In hedges, and bushy grounds : not uncommon.

It flowers in May.

BUGLOSSA sylvestris minor. *Raii Syn.* 227.

ASPERUGO [*procumbens*] *Hudsoni Fl.* 69.

Classis Linnæi Pentandria Monogynia.

Small wild Bugloss. Great Goose-grass. German Madwort.

In corn fields, and among rubbish occasionally: but not very common.

It flowers in April or May.

BUGLOSSUM arvense annuum Lithospermi folio.

Raii Syn. 227.

LITHOSPERMUM [*arvense*] feminibus rugosis, corollis vix calycem superantibus. *Hudsoni Fl.* 66.

Classis Linnæi Pentandria Monogynia.

Bastard Alkanet.

In fields and among corn: common.

It flowers in May or June.

BUGULA *Raii Syn.* 245.

AJUGA [*reptans*] stolonibus reptantibus. *Hudsoni Fl.* 219.

Classis Linnæi Didynamia Gymnospermia.

Bugle.

In woods and moist meadows: common. Found with a white flower occasionally: but uncommon.

It flowers in May.

BULBOCASTANUM. *Raii Syn.* 209.

BUNIAM [Bulbocastanum] *Hudsoni Fl.* 105.

Classis Linnæi Pentandria. Digynia.

Earth-nut. Kipper-nut. Hawk-nut, or rather Hog-nut.

In sandy meadows, and on the Forest between Woodford Row and Chingford Hatch, in plenty.

It flowers in May or June.

BURSA PASTORIS. *Raii Syn.* 306.

THLASPI [Bursa Pastoris] *filiculis obcordatis, foliis radicalibus pinnatifidis. Hudsoni Fl.* 247.

Classis Linnæi Tetradynamia Siliculosa.

Shepherd's Purse.

In fields; on cultivated ground, by the road side; on rubbish, and on old walls: common.

It flowers in April, May, and all the summer months.

BUTOMUS. *Raii Syn.* 273.

BUTOMUS [umbellatus] *Hudsoni Fl.* 152.

Classis Linnæi Enneandria. Hexagynia.

The flowering Rush, or Water Gladiole.

In moist ditches and stagnated waters. Found in many parts of the river Rhodon, in particular between Rey House, Sir James Wright's, and Luxborough House, Sir Edward Walpole's.

It flowers in June.

BUXUS,

WOODFORDIENSES. 21

BUXUS. *Raii Syn.* 445.

BUXUS [*sempervirens*] *Hudsoni Fl.* 355.

Classis Linnæi Monoecia Tetrandia.

The Box Tree.

*Found occasionally near houses, plantations,
and in very old hedges: not very common.*

It flowers in April.

C.

CALAMINTHA humilior folio rotundiore. *Raii*
Syn. 243.

GLECHOMA [*hederacea*] foliis reniformibus crenatis. *Hudsoni Fl.* 224.

Classis Linnæi. Didynamia Gymnospermia.

Ground-Ivy. Gill-go-by-the-grouud. Ale-hoof.
 Tun-hoof.

In shady places, under hedges : very common.
 It flowers in May or June.

CAMPANULA, arvensis erecta, vel speculum Veneris. *Raii Syn.* 278.

CAMPANULA [*hybrida*] caule basi subramoso stricto, foliis oblongis crenatis, calycibus aggregatis, corolla longioribus, capsulis prismaticis. *Hudsoni Fl.* 82.

Classis Linnæi Pentandria Monogynia.

The lesser Venus looking-glass; or, coddled
 Corn Violet.

In fields among corn : Found in a corn field between Ribton Lane and Woodford Bridge : not very common.

It flowers in July or August.

CAMPA-

WOODFORDIENSES. 23

CAMPANULA pratensis flore conglomerato.
Raii Syn. 277.

CAMPANULA [*glomerata*] caule angulato simplici floribus sessilibus, capitulo terminali.
Hudsoni Fl. 82.

Classis Linnæi Pentandria Monogynia.

Little, or, Lesser Throatwort. Canterbury Bells.

In fields on a dry soil: Found in a field near Muncombe: not very common.

It flowers in July.

CAMPANULA rotundifolia. *Raii Syn.* 277.

CAMPANULA [*rotundifolia*] foliis radicalibus reniformibus caulinis linearibus. *Hudsoni Fl.* 80.

Classis Linnæi Pentandria Monogynia.

The lesser round leaved Bell-flower.

In woods and dry barren meadows: very common. On the Forest, every where.

It flowers in August or September.

CAPRIFOLIUM Germanicum. *Raii Syn.* 458.

LONICERA [*Periclymenum*] capitulis ovatis imbricatis terminalibus, foliis omnibus distinctis.
Hudsoni Fl. 80.

Classis Linnæi Pentandria Monogynia.

Common Honeyfuckle or Woodbine.

In hedges: very common, every where.

It flowers from April to August.

CARDAMINE. *Raii Syn.* 299.

CARDAMINE [*pratense*] foliis pinnatis : foliolis radicalibus subrotundis ; caulinis lanceolatis. *Hudsoni Fl.* 256.

Classis Linnæi Tetradynamia Siliquosa.

Common Ladies smock, or Cuckow Flower.

In moist meadows : very common every where.

It flowers in April.

CARDUUS caule crispo. *Raii Syn.* 194.

CARDUUS [*crispus*] foliis sinuatis discurrentibus, margine spinosis, floribus aggregatis terminalibus muticis. *Hudsoni Fl.* 306.

Classis Linnæi Syngenesia : Polygamia Æqualis.

Thistle upon Thistle.

By the road side ; and on dry banks : common.

It flowers in June.

CARDUUS lanceolatus. *Raii Syn.* 195.

CARDUUS [*lanceolatus*] foliis decurrentibus, hastato-pinnatis hispidis : laciniis divaricatis, calycibus spinosis, villosis, caule piloso. *Hudsoni Fl.* 305.

Classis Linnæi Syngenesia : Polygamia Æqualis.
Spear-thistle.

In uncultivated ground, and under hedges : very common.

It flowers in July.

WOODFORDIENSES. 25

CARDUUS Mariæ. *Raii Syn.* 195.

CARDUUS [*Marianus*] foliis amplexi-caulibus hastato-pinnatifidis spinosis, calycibus aphyllis: spinis canaliculatis duplicato-spinosis. *Hudsoni Fl.* 307.

Classis Linnæi Syngenefia : Polygamia Æqualis.
Milk Thistle, or Lady's Thistle.

Found on a bank in the lane leading from Chingford-Hatch to Mr. Lloyd's, Friday Hill House: and in Mr. Warner's Garden, growing spontaneously: not very common.

It flowers in July.

CARDUUS palustris. *Raii Syn.* 194.

CARDUUS [*palustris*] foliis linearibus dentatis decurrentibus: margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Hudsoni Fl.* 306.

Classis Linnæi Syngenefia : Polygamia Æqualis.
Marsh Thistle.

In marshy and shady ground: very common.

It flowers in July.

CARDUUS spinosissimus capitulis minoribus.
Raii Syn. 194.

CARDUUS [*acanthoides*] foliis sinuatis decurrentibus: margine spinosis, calycibus pedunculatis, solitariis erectis villosis. *Hudsoni Fl.* 306.

Classis Linnæi Syngenefia : Polygamia Æqualis.
Wetted Thistle, with small flowers.

Near the road side, on the sides of dry ditches, and on rubbish: not uncommon.

It flowers in June or July.

CARDUUS stellatus. *Raii Syn.* 196.

CENTAUREA [*Calcitrapa*] calycibus subduplicato-squamosis, foliis sessilibus lateralibus pinatifidis linearibus dentatis, caule piloso. *Hudsoni Fl.* 326.

Classis Linnæi Syngenesia : Polygamia Frustranea.

Star Thistle.

In dry fields, and by the road side. Found on the Hill at Woodford Row, near Mr. Warner's large Piece of Water, commonly called, The Gravel Pit : not common.

It flowers in July.

CARDUUS tomentosus, Acanthium dictus, vulgaris. *Raii Syn.* 196.

ONOPORDUM [*Acanthium*] calycibus squarrosis, squamis subulatis, foliis ovato oblongis sinuatis. *Hudsoni Fl.* 308.

Classis Linnæi Syngenesia : Polygamia Æqualis. Cotton Thistle.

By the road side, on rubbish, and, in uncultivated places : common.

It flowers in July.

CARDUUS vulgatissimus viarum. *Raii Syn.* 194.

SERRATULA [*arvensis*] foliis dentatis spinosis. *Hudsoni Fl.* 305.

Classis Linnæi Syngenesia : Polygamia Æqualis.

Common Way Thistle, or creeping Thistle : every where very common.

It flowers in July.

WOODFORDIENSES. 27

CARLINA, sylvestris quibusdam, aliis Atractylis.

Raii Syn. 175.

CARLINA [*vulgaris*] caule multifloro corymbofo, floribus terminalibus. *Hudsoni Fl.* 309.

Classis Linnæi Syngenesia: Polygamia Æqualis.

The common wild Carline Thistle.

In dry Fields, and on barren, sandy soil: common.

It flowers in May.

CARYOPHYLLATA. *Raii Syn.* 253.

GEUM [*urbanum*] floribus erectis, fructu globofo, aristis uncinatis nudis, foliis lyratis.

Hudsoni Fl. 198.

Classis Linnæi Icosandria Polygynia.

Herb-Bennet, or common Avena.

In woods, and under hedges: common.

It flowers in June or July.

CARYOPHYLLUS holosteus arvensis glaber, flore majore. *Raii Syn.* 346.

STELLARIA [*Holostea*] foliis lanceolatis serratis, petalis bifidis. *Hudsoni Fl.* 166.

Classis Linnæi Decandria Trigynia.

The greater Stitchwort.

Under hedges, and on the sides of fields: common.

It flowers in April or May.

CARYOPHYLLUS holosteus arvensis glaber,
flore minore. *Raii Syn.* 346.

STELLARIA [*graminea*] foliis linearibus integerrimis, floribus paniculatis. *Hudsoni Fl.* 166.

Classis Linnæi Decandria Trigynia.

The lesser Stitchwort.

Under hedges, and on the sides of fields, chiefly on a dry soil: common.

It flowers in July.

CARYOPHYLLUS holosteus arvensis medius.
Raii Syn. 347.

STELLARIA [*graminea*] foliis linearibus integerrimis, floribus paniculatis. *Hudsoni Fl.* 166. β.

Classis Linnæi Decandria Trigynia.

The lesser Stitchwort, a variety.

Found sometimes in moist meadows: near Sir Edward Walpole's, Luxborough House, in particular: not common.

It flowers in July.

WOODFORDIENSES. 29

CARYOPHYLLUS latifolius barbatus minor annuus, flore minore. *Raii Syn.* 337.

DIANTHUS [*Armeria*] floribus aggregatis fasciculatis, squamis calycinis lanceolatis villosis tubum æquantibus. *Hudsoni Fl.* 161.

Classis Linnæi Decandria Digynia.

Deptford Pink.

In dry, gravelly, pasture land. Found in the upper end of Sakes, commonly called Snakes-lane: in the lane leading from Chingford Hatch to Mr. Lloyd's, Friday Hill House: and in a field between the end of Horn Lane, and Woodford-Bridge road.

It flowers in July.

CASSIDA palustris vulgatiores flore cæruleo. *Raii Syn.* 244.

SCUTELLARIA [*galericulata*] foliis cordato-lanceolatis rugosis, crenatis, floribus axillaribus. *Hudsoni Fl.* 231.

Classis Linnæi Didynamia Gymnospermia.

Hooded willow herb.

In moist places, near rivers and wet ditches: Found in the meadows, near the river Rhodon: common.

It flowers in July or August.

CASTANEA. *Raii Syn.* 440.

FAGUS [*Castanea*] foliis lanceolatis acuminato-ferratis subtus nudis. *Hudsoni Fl.* 359.

Classis Linnæi Monoecia Polyandria.

The Chesnut Tree.

On the Forest near Wanstead House, Earl Tylney's: not common.

It flowers in May.

CATANANCE leguminosa quorundam. *Raii Syn.*
325.

LATHYRUS [*Nissolia*] pedunculis unifloris,
foliis simplicibus, stipulis subulatis. *Hudsoni*
Fl. 275.

Classis Linnæi Diadelphia Decandria.

Crimson Grass Vetch.

On the sides of fields: Found near Milkwell Field, between the end of Horn Lane, and the Woodford Bridge road, and in a field near Chingford Church: not very uncommon.

It flowers in May, June, or July.

CAUCALIS minor flosculis rubentibn. *Raii Syn.*
219.

CAUCALIS [*Anthriscus*] umbella conferta, seminibus globosis, foliolis ovatis pinnatifidis.
Hudsoni Fl. 99.

Classis Linnæi Pentandria Digynia.

Hedge Parsley.

In bushy grounds, and under hedges.

It flowers in July or August.

CAUCALIS fegetum minor Anthriscu hispido similis. *Raii Syn.* 220.

CAUCALIS [*arvensis*] umbella conferta, involucro universali monophyllo, seminibus oblongis, foliolis ovato-lanceolatis pinnatifidis,
Hudsoni Fl. 98.

Classis Linnæi Pentandria Digynia.

Small Corn Parsley.

In fields and among corn: very common.

It flowers in August.

WOODFORDIENSES. 31

CAUCALIS nodosa, echinato femine. *Raii Syn.*
220.

TORDYLIUM [*nodosum*] umbellis simplicibus,
sessilibus, feminibus exterioribus hispidis.
Hudsoni Fl. 98.

Classis Linnæi Pentandria Digynia.

Knotted Parsley.

*On the sides of fields, and on heaps of loose
stones : not very common.*

It flowers in May.

CENTAURIUM minus. *Raii Syn.* 286.

GENTIANA [*Centaureum*] corollis quinquifidis
infundibuli-formibus caule dichotimo. *Hud-
soni Fl.* 88.

Classis Linnæi Pentandria. Digynia.

Lesser Centaury.

In dry, barren, pasture land : not uncommon.

It flowers from July to October.

CERASUS avium nigra et racemosa. *Raii Syn.*
463.

PRUNUS [*Padus*] floribus racemosis, foliis deci-
duis, basi subtus biglandulosis. *Hudsoni Fl.*
186.

Classis Linnæi Icosandria. Monogynia.

Wild Cluster Cherry, or Bird-Cherry.

*In woods and hedges : Found in a Lane, lead-
ing to Hackney-marsh, near Temple Mills ; and,
between the London Road and West-Ham
Abbey, turning off near the Three Mile Stone :
not common.*

It flowers in May.

CERAS-

CERASTIUM, hirsutum minus, parvo flore ;
Raii Syn. 348.

CERASTIUM [*semidecandrium*] floribus pentandris, petalis emarginatis. *Hudsoni Fl.*
 175.

Classis Linnæi Decandria Pentagynia.

The least mouse-ear Chickweed.

On walls, and gravelly, dry soil : common.

It flowers in April.

CEREFOLIUM sylvestre. *Raii Syn.* 207.

CHÆROPHYLLUM [*temulum*] caule scabro, geniculis tumidis, umbellis cernuis. *Hudsoni Fl.*
 108.

Classis Linnæi Pentandria Digynia.

Wild Chervil.

Under hedges : very common.

It flowers in July or August.

CHAMÆMELUM foetidum. *Raii Syn.* 185.

ANTHEMIS [*Cotula*] receptaculis conicis, paleis fetaceis, feminibus nudis. *Hudsoni Fl.* 323.

Classis Linnæi. Syngenesia : Polygamia superflua.

Stinking May-weed.

In fields, and by the road side : very common.

It flowers in June or July.

CHAMÆMELUM inodorum. *Raii Syn.* 185.

MATRICARIA [*inodora*] receptaculis hemisphæricis, radiis patentibus, feminibus coronato-marginatis, squamis calycinis margine exsoletis. *Hudsoni Fl.* 322.

Classis Linnæi Syngenesia: Polygamia Superflua.
Field Feverfew.

Found on dry gravelly soils; and in barren land: non uncommon.

It flowers in August.

CHAMÆMELUM odoratissimum repens, flore simplici. *Raii Syn.* 185.

ANTHEMIS [*nobilis*] foliis pinnato-compositis linearibus acutis subvillosis. *Hudsoni Fl.* 323.

Classis Linnæi Syngenesia: Polygamia Superflua.
Sweet scented creeping Camomile.

In moist places on the Forest; near Mr. Warner's large Piece of Water, commonly called, The Gravel Pit: common.

It flowers in July or August.

CHAMÆMELUM vulgare. *Raii Syn.* 184.

MATRICARIA [*Chamomilla*] receptaculis conicis, radiis patentibus, feminibus nudis, squamis calycinis margine æqualibus. *Hudsoni Fl.* 322.

Classis Linnæi Syngenesia: Polygamia Superflua.
Corn Feverfew.

In cultivated places; and among corn.

It flowers in June.

CHELEDONIUM minus. *Raii Syn.* 246.

FICARIA [*verna*] *Hudsoni Fl.* 214.

Classis Linnæi Polyandria Polygynia.

Pile-wort, or lesser Celandine.

In meadows, and pasture grounds : very common.

It flowers in April or May.

CHENOPODIUM Betæ folio. *Raii Syn.* 157.

CHENOPODIUM [*polyspermum*] foliis ovatis integerrimis, caule decumbente cymis dichotomis. *Hudsoni Fl.* 92.

Classis Linnæi Pentandria Digynia.

Upright, round-leaved Blite, or All-seed.

Found in cultivated ground, and on dunghills : not very common.

It flowers in July or August.

CHENOPODIUM erectum foliis triangularibus, dentatis, spicis e foliorum alis plurimis longis erectis tenuibus. *Raii Syn.* 155.

CHENOPODIUM [*urbicum*] foliis triangularibus subdentatis, racemis confertis strictissimis caule approximatis longissimis. *Hudsoni Fl.* 89.

Classis Linnæi Pentandria Digynia.

Upright Blite.

On dunghills and ditches by the road side : not uncommon.

It flowers in August or September.

WOODFORDIENSES. 35

CHENOPODIUM Stramonii folio. Chenopodio affinis, folio lato, laciniato, in longissimum mucronem procurrente, florum ramulis sparsis. *Raii Syn.* 154.

CHENOPODIUM [*hybridum*] foliis cordatis, angulo-acuminatis, racemis ramosis nudis. *Hudsoni Fl.* 90.

Classis Linnæi Pentandria Digynia.

Maple leaved Blite.

Found on rich soils, and on dunghills occasionally: not common.

It flowers in August.

CHRYSANTHEMUM segetum. *Raii Syn.* 182.

CHRYSANTHEMUM [*segetum*] foliis amplexicaulibus superne laciniatis, infernè dentato ferratis. *Hudsoni Fl.* 321.

Classis Linnæi Syngenesia: Polygamia Superflua.
Corn Marigold.

In fields, and pasture grounds; among corn in particular; occasionally, on the road side in lanes: not uncommon.

It flowers in June or July.

CICHOREUM sylvestre. *Raii Syn.* 172.

CICHOREUM [*Intybus*] caule simplici racemoso, floribus confertis; omnibus sessilibus. *Hudsoni Fl.* 303.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Wild Succory.

Found on uncultivated places by the road side, and on the sides of fields, chiefly on dry banks: not uncommon.

It flowers in July or August.

CICUTA. *Raii Syn.* 215.

CONIUM [*maculatum*] *feminibus striatis. Hudsoni Fl.* 100.

Classis Linnæi Pentandria Digynia.
Hemlock.

Near Towns and Villages : in shady places by the sides of ditches, and under hedges : not uncommon.

It flowers in June or July.

CICUTARIA tenuifolia. *Raii Syn.* 215.

AETHUSA [*Cynapium*] *Hudsoni Fl.* 107.

Classis Linnæi Pentandria Digynia.

The Lesser Hemlock, or Fool's Parsley.

Found in rich cultivated grounds, and on dung-hills : not uncommon.

It flowers in August or September.

CICUTARIA vulgaris. *Raii Syn.* 207.

CHÆROPHYLLUM [*sylvestre*] *flosculis omnibus fertilibus, caule æquali. Hudsoni Fl.* 108.

Classis Linnæi Pentandria Digynia.

Wild Cicely, or Cow-weed.

Under hedges, in orchards and fields : very common.

It flowers in May or June.

CIRCÆA Lutetiana. *Raii Syn.* 289.

CIRCÆA [*Lutesiana*] caule erecto, racemis pluribus. *Hudsoni Fl.* 9.

Classis Linnæi Diandria Monogynia.

Enchanter's Nightshade.

In shady moist places, under hedges. Found in Ribton Lane and Horn Lane, and in the Copse, the corner of Mr. Warner's large Field: not uncommon.

It flowers in June or July.

CLEMATIS daphnoides major. *Raii Syn.* 268.

VINCA [*major*] caulibus erectis, foliis ovatis. *Hudsoni Fl.* 77.

Classis Linnæi Pentandria Monogynia.

The greater Periwinkle.

In woods and under hedges. Found under the hedge which parts Mr. Warner's Field from Ribton-Lane: and on the Islands in Mr. Warner's large Piece of Water, on the Hill before his House, commonly called, The Gravel Pit: not common.

It flowers in May.

CLEMATIS latifolia; seu Atragene quibusdam. *Raii Syn.* 258.

CLEMATIS [*Vitalba*] foliis primatis; foliolis cordatis scandentibus. *Hudsoni Fl.* 217.

Classis Linnæi Polyandria Polygynia.

Great wild Climber. Traveller's Joy. Bethwyne.

In Hedges. Found in Hagger-Lane, leading from the Forest near the Wind-Mill, to Walthamstow: and on the side of the road between Chigwell and Abridge: not very common.

It flowers in July.

CLINOPODIUM Origano simile. *Raii Syn.*

239.

CLINOPODIUM [*vulgare*] capitulis subrotundis hispidis, bracteis fetaceis. *Hudsoni Fl.* 228.

Classis Linnæi Didynamia Gymnospermia.

Great wild Basil.

Found sometimes under hedges, overgrown with brambles : not common.

It flowers in July.

CONVOLVULUS major. *Raii Syn.* 275.

CONVOLVULUS [*sepium*] foliis sagittatis postice truncatis, pedunculis tetragonis unifloris.

Hudsoni Fl. 74.

Classis Linnæi Pentandria Monogynia.

Great Bindweed.

Found under hedges ; particularly in moist places : very common.

It flowers in July or August.

CONVOLVULUS minor vulgaris. *Raii Syn.* 275.

CONVOLVULUS [*arvensis*] foliis sagittatis utrinque acutis, pedunculis unifloris. *Hudsoni Fl.*

74.

Classis Linnæi Pentandria Monogynia.

Small Bindweed.

In fields, and by the road side : common.

It flowers in June or July.

WOODFORDIENSES. 39

CONYZA media. *Raii Syn.* 174.

INULA [*dysenterica*] foliis oblongis, caule hirsuto peniculato, squamis calycinis fetaceis.

Hudsoni Fl. 320.

Classis Linnæi Syngenesia: Polygamia Superflua.
Middle Fleabane.

Found in moist places, on the banks of rivers, and near moist ditches: common.

It flowers in August.

CONYZA minor. *Raii Syn.* 174.

INULA [*Pulicaria*] foliis undulatis, amplexicaulibus, caule prostrato. *Hudsoni Fl.* 320.

Classis Linnæi Syngenesia: Polygamia Superflua.
Small Fleabane.

Found in places where there have been stagnated waters in winter, and in other low places: not uncommon.

It flowers in August or September.

CORNUS foemina. *Raii Syn.* 460.

CORNUS [*sanguinea*] arborea, cymis nudis.
Hudsoni Fl. 58.

Classis Linnæi Tetrandia Monogynia.

The Female Cornel. Dog-berry tree. Gatterwood, or Prick-wood.

Found in woods; and in hedges by the road side: common.

It flowers in June.

CORYLUS sylvestris. *Raii Syn.* 439.

CORYLUS [*Avellana*] stipulis ovatis obtusis.
Hudsoni Fl. 360.

Classis Linnæi Monoecia Polyandria.

The Hazel-nut Tree.

*Found upon the woody Parts of the Forest, and
in hedges : common.*

It flowers in March.

CRACCA. *Raii Syn.* 322.

VICIA [*Cracca*] pedunculis multifloris, floribus
imbricatis, foliolis lanceolatis pubescentibus,
stipulis integris. *Hudsoni Fl.* 277.

Classis Linnæi Diadelphia Decandria.

Tufted Vetch.

Found under hedges : not uncommon.

It flowers in July or August.

CRACCA minor. *Raii Syn.* 322.

ERVUM [*hirsutum*] pedunculis multifloris, semi-
nibus globosis binis. *Hudsoni Fl.* 280.

Classis Linnæi Diadelphia Decandria.

Small wild Tare. Tine Tare, or Hairy Tare.

*Found in fields and under hedges : very com-
mon.*

It flowers in July.

WOODFORDIENSES. 41

CRACCA minor filiquis singularibus, flosculis cærulescentibus. *Raii Syn.* 322.

ERVUM [*tetraspermum*] pedunculis sub-bifloris feminibus globosis quaternis. *Hudsoni Fl.* 280.

Classis Linnæi Diadelphia Decandria.

Tine Tare with small pods, or smooth Tare.

Found in fields, and under hedges: very common.

It flowers in June.

CUSCUTA major. *Raii Syn.* 281.

CUSCUTA [*Europæa*] floribus sessilibus. *Hudsoni Fl.* 89.

Classis Linnæi Pentandria Digynia.

Dodder. Hell-weed. Devil's Guts.

On forests, and among leguminous plants. Found on the Forest, near The Eagle at Snarebrooke, on the south-west side of the great pond, climbing round the Erica Vulgaris: not common.

It flowers in July.

CYANUS. *Raii Syn.* 198.

CENTAUREA [*Cyanus*] calycibus ferratis, foliis linearibus integerrimis; infimis dentatis. *Hudsoni Fl.* 325.

Classis Linnæi Syngenesia: Polygamia Frustranea.

Blue Bottles, or Corn Bottles.

In fields, and particularly among corn: common.

It flowers in July.

CYNOGLOSSA folio virenti. *Raii Syn.* 226.

CYNOGLOSSUM [*officinale*] staminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus. *Hudsoni Fl.* 67. β.

Classis Linnæi Pentandria Monogynia.

The lesser green leaved Hounds-tongue.

In uncultivated places, on rubbish, and by the sides of Lanes : Found on the side of the road near Mr. Lloyd's, Friday-Hill House, and in the road from London to Woodford-bridge, between the six and seven Mile Stone, near Mobs-hole : not common.

It flowers in June.

CYNOGLOSSUM. *Raii Syn.* 226.

CYNOGLOSSUM [*officinale*] staminibus corolla brevioribus, foliis lato-lanceolatis, tomentosis sessilibus. *Hudsoni Fl.* 67.

Classis Linnæi Pentandria Monogynia.

Great Hounds-tongue.

In uncultivated places, on rubbish, and by the road side : not uncommon.

It flowers in June.

D.

DAMASONIUM stellatum Dalechampii. *Raii*
Syn. 272.

ALISMA [*Damasonium*] foliis cordato-oblongis
floribus hexaginis. *Hudsoni Fl.* 138.

Classis Linnæi. Hexandria Polygynia.

Star-headed Water Plantain.

Found in a Pond near Ilford, and in Mr. Warner's large Pond, commonly called The Gravel Pit, on the Hill, opposite Hart's, his House in Woodford Row : near London, common : but not elsewhere.

It flowers from June to August.

DAUCUS VULGARIS. *Raii Syn.* 218.

DAUCUS [*Carota*] seminibus hispidis. *Hudsoni*
Fl. 99.

Classis Linnæi Pentandria Digynia.

Wild Carrot, or Bird's-nest.

In meadows and pastures : very common.

It flowers in June or July.

DELPHINIUM fegetum flore cæruleo. *Raii Syn.* 273.

DELPHINIUM [*Consolida*] nectariis monophyllis, caule subdiviso. *Hudsoni Fl.* 207.

Classis Linnæi Polyandria Trigynia.

Wild Larkspur.

Found in a little inclosure between Captain Hunt's, Woodford Hall, and the road: not common.

It flowers in June.

DENS LEONIS. *Raii Syn.* 170.

LEONTODON [*Taraxacum*] calyce inferne reflexo. *Hudsoni Fl.* 297.

Classis Linnæi Syngenesia: Polygamia Æqualis. Dandelion.

In meadows and pastures: very common every where, and in cultivated ground, in particular.

It flowers from April to July.

DENS LEONIS hirsutus λεπτόκαυλον, Hieracium dictus. *Raii Syn.* 171.

LEONTODON [*hispidum*] calyce toto erecto hispido, foliis hispidis dentatis integerrimis; fetis furcatis. *Hudsoni Fl.* 297.

Classis Linnæi Syngenesia: Polygamia Æqualis.

Rough Dandelion, commonly called Dandelion Hawkweed.

Found in meadows and pastures: common.

It flowers in May, June, or July.

DIGITALIS purpurea. *Raii Syn.* * 283.

DIGITALIS [*purpurea*] calycinis foliolis ovatis acutis, corollis obtusis labio superiore integro.

Hudsoni Fl. 240.

Classis Linnæi Didynamia Angiospermia.

Purple Fox-glove.

By the road side, under hedges, near old gravel-pits, and in the shady parts of the Forest: common. It is sometimes found with a white flower; but very uncommon.

It flowers in July or August.

DIPSACUS minor, seu virga pastoris. *Raii Syn.* 192.

DIPSACUS [*pilosus*] foliis petiolatis, appendiculatis. *Hudsoni Fl.* 49.

Classis Linnæi Tetrandia Monogynia.

Small Wild Teasel, or Shepherd's Rod.

In moist places, and under hedges: Found near Mr. Lloyd's, Friday Hill House, in the Lane leading to Chingford Church: not common.

It flowers in July or August.

DIPSACUS sylvestris, seu Labium Veneris. *Raii Syn.* 192.

DIPSACUS [*sylvestris*] foliis sessilibus serratis, aristis fructus erectis. *Hudsoni Fl.* 49.

Classis Linnæi Tetrandria Monogynia.

Wild Teasel.

In uncultivated moist places, and under hedges: very common.

It flowers in July.

E.

ECHIUM vulgare. *Raii Syn.* 227.

ECHIUM [*vulgare*] caule simplici erecto, foliis caulinis lanceolatis hispidis, floribus spicatis lateralibus staminibus corollam æquantibus.

Hudsoni Fl. 69.

Classis Linnæi Pentandria Monogynia.

Viper's Bugloss.

By the road side, on dry barren land, and sometimes on walls : not very common.

It flowers in July.

EQUISETUM arvense et palustre longioribus setis. *Raii Syn.* 130.

EQUISETUM [*arvense*] scapo fructificante nudo sterili frondoso. *Hudsoni Fl.* 380.

Classis Linnæi Cryptogamia : Filices.

Corn Horsetail.

In moist fields, and among corn.

It flowers in March.

EQUI-

EQUISETUM palustre. *Raii Syn.* 131.

EQUISETUM [*palustre*] caule angulato, frondibus simplicibus. *Hudsoni Fl.* 380.

Classis Linnæi Cryptogamia : Filices.

The Lesser Marsh Horse-tail.

In Fields in moist places.

It flowers in June.

EQUISETUM sylvaticum. *Raii Syn.* 130.

EQUISETUM [*sylvaticum*] caule spicato, frondibus compositis. *Hudsoni Fl.* 380.

Classis Linnæi Cryptogamia : Filices.

Wood Horse-tail.

In shady moist places in woods, and sometimes on the banks of rivers : not very common.

It flowers in April or May.

ERICA Brabantica folio coridis hirsuto quaternis. *Raii Syn.* 471.

ERICA [*Tetralix*] antheris bicornibus inclusis, corollis subglobosis aggregatis calyce longioribus foliis quaternis ciliatis patentibus. *Hudsoni Fl.* 144.

Classis Linnæi Octandria Monogynia.

Cross-leaved Heath.

Found near the great Pond at Snaresbrook, on the Forest : not very common.

It flowers in July, August, and September.

ERICA tenuifolia. *Raii Syn.* 471.

ERICA [*cinerea*] antheris bicornibus inclusis, corollis ovatis racemosis, foliis ternis glabris linearibus. *Hudsoni Fl.* 144.

Classis Linnæi Octandria Monogynia.

Fine-leaved Heath.

Found on the Forest, near the great Pond at Snaresbrook; and not far from The Windmill: not very common.

It flowers from June to August.

ERICA vulgaris. *Raii Syn.* 470.

ERICA [*vulgaris*] antheris bicornibus inclusis corollis inæqualibus campanulatis mediocribus, foliis oppositis sagittatis. *Hudsoni Fl.* 144.

Classis Linnæi Octandria Monogynia.

Common Heath or Ling.

On the Forest: not uncommon.

It flowers from June to September.

ERUCA hirsuta, filiqua cauli appressa, Erysimum dicta. Erysimum vulgare. *Raii Syn.* 298.

ERYSIMUM [*officinale*] filiquis spicæ appressis. *Hudsoni Fl.* 250.

Classis Linnæi. Tetradynamia Siliquosa.

Hedge mustard.

Under hedges, by the road side, and among rubbish: very common.

It flowers in May.

ERUCA lutea, seu Barbarea *Raii Syn.* 297.

ERYSIMUM [*Barbarea*] foliis lyratis: extimo subrotundo. *Hudsoni Fl.* 251.

Classis Linnæi Tetradynamia Siliquosa.

Winter Cresses, or Rocket.

In ditches, and moist places: very common: in cultivated ground: not uncommon.

It flowers in May.

ERYSIMUM Sophia dictum. Sophia Chirurgorum. *Raii Syn.* 298.

SISYMBRIUM [*Sophia*] petalis calyce minoribus, foliis decomposito pinnatis. *Hudsoni Fl.* 259.

Classis Linnæi. Tetradynamia Siliquosa.

Flix-weed.

Among rubbish, on old walls, and sometimes under hedges: not uncommon.

It flowers in June.

EUONYMUS vulgaris, *Raii Syn.* 468.

EUONYMUS [*Europæus*] floribus plerisque quadrifidis. *Hudsoni Fl.* 84.

Classis Linnæi Pentandria Monogynia.

Spindle Tree, or Prick-wood.

In woods and hedges: Found in the Woodford-bridge Road, near the eight mile stone: not uncommon.

It flowers in April and May.

EUPATORIUM Cannabinum. *Raii Syn.* 179.

EUPATORIUM [*Cannabinum*] foliis digitatis.
Hudsoni Fl. 310.

Classis Linnæi Syngenesia : Polygamia Æqualis.

Common Hemp Agrimony, or Dutch Agrimony:
Found by the Sides of the River Rhodon : common.

It flowers in July or August.

EUPHRASIA. *Raii Syn.* * 284.

EUPHRASIA [*officinalis*] foliis ovatis argute
dentatis. *Hudsoni Fl.* 234.

Classis Linnæi Didynamia Angiospermia.

Eyebright.

In every part of the Forest : common.

It flowers in August or September.

EUPHRASIA pratensis rubra. *Raii Syn.* * 284.

EUPHRASIA [*odontites*] foliis linearibus : omnibus
ferratis. *Hudsoni Fl.* 234.

Classis Linnæi Didynamia Angiospermia.

Red Eyebright.

In fields and pasture grounds : common.

It flowers in July or August.

F

FAGUS. *Raii Syn.* 439.

FAGUS [*sylvatica*] foliis ovatis obsolete ferratis.
Hudsoni Fl. 360.

Classis Linnæi Monoecia Polyandria.

The Beech Tree.

In different Parts of the Forest: not uncommon.

It flowers in May.

PEGOPYRUM scandens sylvestre. *Raii Syn.*
144.

POLYGONUM [*Convolutus*] foliis cordatis caule
volubili, floribus planiusculis. *Hudsoni Fl.*
149.

Classis Linnæi Octandria Trigynia.

Black Bindweed.

*Among corn, in gardens, and cultivated places:
not uncommon.*

It flowers from June to September.

FILIPENDULA. *Raii Syn.* 259.

SPIRÆA [*Filipendula*] foliis pinnatis : foliolis uniformibus ferratis caule herbaceo, floribus cymosis. *Hudsoni Fl.* 190.

Classis Linnæi Icosandria Pentagynia.

Common Dropwort.

Found on a billy field near Chingford Church : not common.

It flowers in July.

FILIX foemina. *Raii Syn.* 124.

PETERIS [*aquilina*] fronde supradecomposita : foliolis pinnatis : pinnis lanceolatis : infimis pinnatifidis ; superioribus minoribus. *Hudsoni Fl.* 384.

Classis Linnæi Cryptogamia : Filices.

Female Fern, or common Brakes.

In woods ; and on the Forest, every where.

It flowers in August.

FILIX mas non ramosa, pinnulis latis auriculatis spinosis. *Raii Syn.* 121.

POLYPODIUM [*aculeatum*] fronde bipinnata : pinnis lanceolato-ovatis dentato-ciliatis, stipite strigoso. *Hudsoni Fl.* 389.

Classis Linnæi Cryptogamia.

Prickly Auriculate Male Fern. Female Polypody.

On banks, and the shady side of ditches : not very common. Found under a hedge in Ribton-Lane, and under a hedge the side of Mrs. North's wood, in the road from Woodford Row to Muncombe.

WOODFORDIENSES. 53

FILIX mas ramosa, pinnulis dentatis. *Raii Syn.*
124.

POLYPODIUM [*cristatum*] frondibus sub pinnatis : foliolis ovato oblongis ; pinnis obtusifusculis apice acute ferratis. *Hudsoni Fl.* 390.

Classis Linnæi Cryptogamia : Filices.

Great branched Fern, with indented leaves ; or, cristed Polypody.

In woods, and moist shady places : not uncommon. Found in the thickets on the sides of the great Pond at Snaresbrook.

FILIX ramosa minor. *Raii Syn.* 125.

POLYPODIUM [*Dryopteris*] fronde supra decomposita : foliolis ternis bipinnatis. *Hudsoni Fl.*
391.

Classis Linnæi Cryptogamia : Filices.

The lesser branched Fern, or branched Polypody.

Found on the walls of Chingford Church : very uncommon.

FILIX ramosa non dentata florida. *Raii Syn.*
125.

OSMUNDA [*regalis*] fronde bipennata apice racemifera. *Hudsoni Fl.* 382.

Classis Linnæi Cryptogamia : Filices.

Water Fern. Flowering Fern, or Osmund Royal.

In moist boggy places : very uncommon ; but found on the side of a gravel-pit, behind Mr. Moxon's House, near the path leading from thence to Fair Mead bottom.

It flowers in July or August.

FILIX

FILIX mas vulgaris. *Raii Syn.* 120.

POLYPODIUM [*Filix mas*] fronde bipinnata: pinnulis obtusis crenulatis, stipite paleaceo.

Hudsoni Fl. 389.

Classis Linnæi Cryptogamia: Filices.

Common Male Fern.

On banks under hedges, chiefly in the shade: very common.

FOENICULUM vulgare. *Raii Syn.* 217.

ANETHUM [*Foeniculum*] fructibus ovatis. *Hudsoni Fl.* 110.

Classis Linnæi Pentandria Digynia.

Common Fennel, or Finckle.

On dry soil. Found in a Lane in Woodford, which leads from Dr. Wilkinfon's, to The Windmill.

It flowers in August.

FRAGARIA. *Raii Syn.* 254.

FRAGARIA [*vesca*] flagellis reptans. *Hudsoni Fl.* 194.

Classis Linnæi Icosandria Polygynia.

Strawberry.

In woods, and in the shady part of the Forest: not uncommon.

It flowers in April or May.

WOODFORDIENSES. 55

FRAGARIA sterilis. *Raii Syn.* 254.

FRAGARIA [*sterilis*] caule decumbente repente. *Hudsoni Fl.* 195.

Classis Linnæi Icofandria Polygynia.

Barren Strawberry.

In barren land, and on the Forest : very common.

It flowers in April.

FRANGULA, five alnus nigra baccifera. *Raii Syn.* 465.

RHAMNUS [*Frangula*] inermis, floribus monogynis hermaphroditis, foliis integerrimis. *Hudsoni Fl.* 83.

Classis Linnæi Pentandria Monogynia.

The Black berry-bearing Alder.

In moist woods, and hedges : Found on the south-west side of the great Pond at Snaresbrook, near the Path, at the head of it : not common.

It flowers in April or May.

FRAXINUS. *Raii Syn.* 469.

FRAXINUS [*excelsior*] foliis ferratis, floribus apetalis. *Hudsoni Fl.* 379.

Classis Linnæi Polygamia Dioecia.

The common Ash Tree.

In hedges, and sometimes in woods : very common.

It flowers in March and April,

FUMARIA vulgaris. *Raii Syn.* 204.

FUMARIA [*officinalis*] pericarpis monospermis
racemosis, caule diffuso. *Hudsoni Fl.* 270.

Classis Linnæi Diadelphia Hexandria.

Fumitory.

*In fields, banks, and sometimes on walls :
common.*

It flowers in April, May, or June.

FUMARIA major scandens flore pallidior. *Raii
Syn.* 204.

FUMARIA [*capreolata*] pericarpis monospermis
racemosis, foliis scandentibus subcirrhosis.
Hudsoni Fl. 270.

Classis Linnæi Diadelphia Hexandria.

The greater ramping Fumitory.

*In fields, banks, and sometimes in woods, and
on walls : not very common.*

It flowers in August.

G.

GALEOPSIS legitima Dioscoridis. *Raii Syn.*
237.

STACHYS [*sylvatica*] verticillis sexfloris, foliis
cordatis petiolatis. *Hudsoni Fl.* 227.

Classis Linnæi Didynamia Gymnospermia.
Hedge Nettle.

In woods, and under hedges: very common.

It flowers in July or August.

GALLIUM luteum. *Raii Syn.* 224.

GALLIUM [*verum*] foliis octonis linearibus sul-
catis, ramis floriferis brevibus. *Hudsoni Fl.*
55.

Classis Linnæi Tetrandia Monogynia.

Yellow Ladies Bedstraw, or Cheese-rening.

*In dry meadows, and on the sides of fields and
little hillocks: very common.*

It flowers in July or August.

GENISTA angulosa trifolia. *Raii Syn.* 474.

SPARTIUM [*scoparium*] foliis ternatis solitariisque, ramis inermibus angulatis. *Hudsoni Fl.* 271.

Classis Linnæi Diadelphia Decandria.

Common Broom.

On the Forest, and in dry fields: very common.

It flowers in May or June.

GENISTA minor Aspalathoides. *Raii Syn.* 475.

GENISTA [*Anglica*] spinis simplicibus, ramis stoloniferis inermibus foliis lanceolatis. *Hudsoni Fl.* 272.

Classis Linnæi Diadelphia Decandria.

Needle Furze or Petty Whin.

On moist parts of the Forest: very common.

It flowers in May or June, sometimes in April.

GENISTA spinosa minor. *Raii Syn.* 475.

ULEX [*Europæus*] foliis villosis acutis, spinis sparsis. *Hudsoni Fl.* 272. β .

Classis Linnæi Diadelphia Decandria.

The lesser Furze, Whins or Gorfe.

On the Forest, and in woods: common.

It flowers in July, August, or September.

GENISTA spinosa vulgaris. *Raii Syn.* 475.

ULEX [*Europæus*] foliis villosis acutis, spinis
sparsis. *Hudsoni Fl.* 272.

Classis Linnæi Diadelphia Decandria.

Furze, Whins, or Gorse.

On the Forest, and in woods: very common.

It flowers in April, May, or June.

GENISTELLA tinctoria. *Raii Syn.* 474.

GENISTA [*tinctoria*] foliis lanceolatis glabris,
ramis striatis teretibus erectis. *Hudsoni Fl.*
272.

Classis Linnæi Diadelphia Decandria.

Wood-waxen, or Dyer's Weed. Green-wood.

*In meadows and fields: very common: when if
cows feed upon it, it gives a bitter taste to their
milk.*

It flowers in July.

GERANIUM cicutæ folio, inodorum. *Raii Syn.*
357.

GERANIUM [*cicutarium*] pedunculis multifloris,
calycibus pentaphyllis, floribus pentandris,
foliis pinnatis incisus obtusis. *Hudsoni Fl.*
262.

Classis Linnæi Monadelphia Decandria.

Field Cranesbill without scent, or, Hemlock
leaved Cranesbill.

*On rubbish, and by the road side: not uncom-
mon.*

It flowers from April to June.

GERANIUM columbinum. *Raii Syn.* 359.

GERANIUM [*molle*] pedunculis bifloris, foliisque floralibus alternis, caule ramofo diffuso, calycibus muticis. *Hudsoni Fl.* 265.

Classis Linnæi Monadelphia Decandria.

Common Dove's foot. Cranesbill.

In fields, meadows, pastures, and under hedges: very common.

It flowers in May or June.

GERANIUM columbinum humile flore cæruleo minimo. *Raii Syn.* 359.

GERANIUM [*pusillum*] pedunculis bifloris, petalis bifidis, caule prostrato, foliis reniformibus palmatis linearibus acutis. *Hudsoni Fl.* 266.

Classis Linnæi Monadelphia Decandria.

Small-flowered Dove's foot. Cranesbill.

In fields and by the road side on a dry, sandy soil: Found on a bank near Leighton: not common.

It flowers in June.

GERANIUM columbinum majus foliis dissectis. *Raii Syn.* 359.

GERANIUM [*dissectum*] pedunculis bifloris, foliis quinque-partito-bifidis, petalis emarginatis longitudine calycis, capsulis villosis. *Hudsoni Fl.* 266.

Classis Linnæi Monadelphia Decandria.

Dove's foot. Cranesbill with jagged leaves.

In fields and meadows, on a dry gravelly soil.

It flowers in May, June, or July.

WOODFORDIENSES. 61

GERANIUM Robertianum. *Raii Syn.* 358.

GERANIUM [*Robertianum*] pedunculis bifloris
calycibus pilosis decemangulatis. *Hudsoni Fl.*
264.

Classis Linnæi Monadelphia Decandria.

Herb. Robert.

In hedges, and on walls and rubbish : common.

It flowers in April, May, or June.

GNAPHALIUM, longifolium humile ramosum,
capitulis nigris nobis. *Raii Syn.* 181.

GNAPHALIUM [*uliginosum*] caule ramoso diffuso,
floribus confertis terminalibus. *Hudsoni Fl.*
313.

Classis Linnæi Syngenesia: Polygamia Superflua.

Black-headed, long-leaved, low, branched Cudweed.

Found in dry ditches, and where there have been stagnated waters in winter : not uncommon.

It flowers in August.

GNAPHALIUM minimum. *Raii Syn.* 181.

FILAGO [*montana*] floribus conicis terminalibus
axillaribusque, caule dichotomo erecto. *Hudsoni Fl.* 328.

Classis Linnæi Syngenesia: Polygamia Necessaria.

The least Cudweed.

In sandy, dry meadows, and often among corn.

It flowers in June or July.

GNAPHALIUM minus, seu herba impia. *Raii*
Syn. 180.

FILAGO [*Germanica*] calycibus pyramidatis
 pentagonis axillaribus, caule dichotomo erectiusculo. *Hudsoni Fl.* 328.

Classis Linnæi Syngenesia: Polygamia Necessaria.

Common Cudweed.

In dry meadows, and by the road side: not uncommon.

It flowers in June or July.

H.

HEDERA communis major et minor. *Raii Syn.* 459.

HEDERA [*Helix*] foliis ovatis lobatisque. *Hudsoni Fl.* 85.

Classis Linnæi Pentandria Monogynia.

Climbing or Berried Ivy : also barren or creeping Ivy.

In woods or hedges : very common : There is now a very flourishing tree of it, against the east end of Chingford Church, 2 feet 7 inches in circumference.

It flowers in October.

HELENIUM. *Raii Syn.* 176.

INULA [*Helenium*] foliis ovatis rugosis ; subtus tomentosis, calycum squamis ovatis. *Hudsoni Fl.* 319.

Classis Linnæi Syngenesia : Polygamia Superflua.

Elecampane.

In meadows and pastures. Found near a farmhouse, on the right hand of the lane leading to Muncombe : not very common.

It flowers in July and August.

HELLEBORASTER maximus. *Raii Syn.* 271.

HELLEBORUS [*foetidus*] caule multifloro folioso,
foliis pedatis. *Hudsoni Fl.* 215.

Classis Linnæi Polyandria Polygynia.

Great Bastard black Hellebore, Bear's Foot or
Setterwort.

*Found opposite a Farm house at Muncombe,
and near Mr. Warner's Gravel Pit Pond: not
common.*

It flowers in February or March.

HESPERIS allium redolens. *Raii Syn.* 293.

ERYSIMUM [*Alliaria*] foliis cordatis. *Hudsoni
Fl.* 251.

Classis Linnæi Tetradynamia Siliquosa.

Jack by the Hedge, or Sawce alone.

*Under hedges, and by the sides of ditches:
very common.*

It flowers in May.

HIERACIUM Echoides capitulis Cardui bene-
dicti. *Raii Syn.* 166.

PICRIS [*Echinoides*] perianthiis involuero maxi-
mo cinctis. *Hudsoni Fl.* 294.

Classis Linnæi Syngenesia: Polygamia Æqualis.

Lang de beuf, or, Ox Tongue.

On the sides of fields: common.

It flowers in July or August.

HIERACIUM fruticosum latifolium hirsutum.

Raii Syn. 167.

HIERACIUM [*sabaudum*] caule erecto multifloro, foliis ovato-lanceolatis dentatis semi-amplexicaulibus. *Hudsoni Fl.* 300.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Bushy Hawkweed with broad rough Leaves.

On heaths, commons, and a dry soil: not uncommon.

It flowers in June or July.

HIERACIUM longius radicum. *Raii Syn.* 165.

HYPOCHÆRIS [*radicata*] foliis dentato sinuatis scabris, caule ramoso nudo, pedunculis squamosis. *Hudsoni Fl.* 302.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Long rooted Hawkweed.

In fields; and in the open parts of the Forest: very common.

It flowers in May, June, or July.

HIERACIUM luteum glabrum, five minus hirsutum. *Raii Syn.* 165.

CREPIS [*tectorum*] foliis lanceolatis sagittatis sessilibus levibus, inferioribus dentatis. *Hudsoni Fl.* 301.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Smooth Succory Hawkweed.

In fields, on old walls, and on the roofs of houses: common.

It flowers from June to September.

HIERACIUM minus præmorfa radice. *Raii Syn.* 164.

LEONTODON [*autumnale*] foliis glabris dentatis integerrimis, calycibus erectis, pedunculis squamosis. *Hudsoni Fl.* 297.

Classis Linnæi Syngenesis: Polygamia Æqualis.
Hawkweed with bitten roots. Yellow Devil's bit.

In fields ; and in the open parts of the Forest : very common.

It flowers in July or August.

HORMINUM sylvestre Lavendulæ flore. *Raii Syn.* 237.

SALVIA [*verbena*] foliis sinuato-ferratis, corollis calyce angustioribus. *Hudsoni Fl.* 9.

Classis Linnæi Diandria Monogynia.

Common English wild Clary : sometimes called Oculus Christi.

In meadows, and on a dry soil. Found often in Church yards ; in that of Layton, commonly called Low Layton, and Woodford ; on account perhaps of its use in Medicine, as mentioned by Ray, and so made use of as an instance of Popish imposture : not very common.

It flowers in May.

HYACINTHUS Anglicus. *Raii Syn.* 373.

HYACINTHUS [*non scriptus*] corollis campanulatis sexpartitis apice revolutis. *Hudsoni Fl.*

123.

Classis Linnæi Hexandria Monogynia.

English Hyacinth or Harebells.

In woods, and under hedges: common. Found in Mr. Warner's large field, in plenty.

It flowers in May.

HYDROCERATOPHYLLON folio aspero, quatuor cornibus armato. *Raii Syn.* 135.

CERATOPHYLLUM [*demersum*]. *Hudsoni Fl.*

357.

Classis Linnæi Monoecia Polyandria.

Horned Pondweed.

In stagnated waters, ditches, and rivers where there is little stream. Found in Mr. Warner's, and most of the Ponds on the Forest: very common.

It flowers in July.

HYDROCOTYLE vulgaris. *Raii Syn.* 222.

HYDROCOTYLE [*vulgaris*] foliis peltatis, umbellis quinquefloris. *Hudsoni Fl.* 96.

Classis Linnæi Pentandria Digynia.

Marsh Pennywort, or White Rot.

In marshes, and boggy ground: very common.

It flowers in May.

HYOSCYAMUS vulgaris. *Raii Syn.* 274.

HYOSCYAMUS [*niger*] foliis amplexicaulibus.
Hudsoni Fl. 77.

Classis Linnæi Pentandria Monogynia.

Common Henbane.

Near villages, by the way-side, in ditches and among rubbish : common.

It flowers in June.

HYPERICUM. *Raii Syn.* 342.

HYPERICUM [*perforatum*] floribus trigynis,
caule ancipiti, foliis obtusis pellucido-punctatis. *Hudsoni Fl.* 290.

Classis Linnæi Polyadelphia Polyandria.

Saint John's Wort.

Under hedges in fields, and by the road side : common.

It flowers in July.

HYPERICUM Androsæmum dictum. *Raii Syn.* 343.

HYPERICUM [*hirsutum*] floribus trigynis, calycibus ferrato-glandulosis, caule erecto, foliis ovatis pubescentibus. *Hudsoni Fl.* 291.

Classis Linnæi Polyadelphia Polyandria.

Park-leaves. Tutshan, or Hairy Saint John's Wort.

In woods, under hedges, and in the sides of fields : common.

It flowers in July.

HYPERICUM Ascyron dictum caule quadrangulo. *Raii Syn.* 344.

HYPERICUM [*quadrangulum*] floribus trigynis, caule quadrato herbaceo. *Hudsoni Fl.* 292.

Classis Linnæi Polyadelphia Polyandria.

Saint Peter's Wort.

Under hedges in moist ground ; and near the river Rhodon : common.

It flowers in July.

HYPERICUM maximum Androsæmum vulgare dictum. *Raii Syn.* 343.

HYPERICUM [*Androsæmum*] floribus trigynis fructibus baccatis, caule fruticoso ancipiti. *Hudsoni Fl.* 291.

Classis Linnæi Polyadelphia Polyandria.

Tutsan or Park-leaves.

In woods, and under hedges : Found in Sakes, commonly called, Snakes Lane : not very common.

It flowers in July.

HYPERICUM minus supinum. *Raii Syn.* 343.

HYPERICUM [*humifusum*] floribus trigynis axillaribus solitariis, caulibus ancipitibus filiformibus, foliis glabris. *Hudsoni Fl.* 290.

Classis Linnæi Polyadelphia Polyandria.

The least trailing Saint John's Wort.

On dry lands, and by the sides of fields : not uncommon.

It flowers in July.

HYPERICUM pulchrum Tragi. *Raii Syn.* 342.

HYPERICUM [*pulchrum*] floribus trigynis, calycibus ferrato-glandulosis, foliis cordatis glabris, bracteis integerrimis, caule tereti. *Hudsoni Fl.* 290.

Classis Linnæi Polyadelphia Polyandria.

Upright Saint John's Wort.

In woods, and under hedges: common.

It flowers in July.

JACEA major. *Raii Syn.* 198.

CENTAUREA [*Scabiosa*] calycibus ciliatis, foliis
pinnatifidis : laciniis lanceolatis. *Hudsoni Fl.*
325.

Classis Linnæi Syngenesia : Polygamia Frustra-
nea.

Great Knap-weed, or Matfellow.

On the side of fields : common.

It flowers in June or July.

JACEA nigra. *Raii Syn.* 198.

CENTAUREA [*Jacea*] calycibus squamosis, foliis
lanceolatis : radicalibus sinuato dentatis, ramis
angulatis. *Hudsoni Fl.* 326.

Classis Linnæi Syngenesia : Polygamia Frustra-
nea.

Common Knap-weed, or Matfellow.

On the side of fields : common.

It flowers in July or August.

JACOBÆA latifolia, palustris five aquatica. *Raii Syn.* 178.

SENECIO [*aquaticus*] corollis radiantibus, foliis dentatis, radicalibus ovatis, caulinis pinnatifidis; lacinia extima majore. *Hudsoni Fl.* 317.

Classis Linnæi Syngenefia: Polygamia Superflua.
Broad leaved Marsh, or Water Ragwort.

In ditches, and moist boggy places: very common.

It flowers in July or August.

JACOBÆA senecionis folio incano perennis. *Raii Syn.* 177.

SENECIO [*erucifolius*] corollis radiantibus, foliis pinnatifidis dentatis incanis, caule erecto. *Hudsoni Fl.* 317.

Classis Linnæi Syngenefia: Polygamia Superflua.
Hoary perennial Ragwort, with Groundsel leaves.

In meadows, woods, and under hedges: common.

It flowers in July.

JACOBÆA vulgaris. *Raii Syn.* 177.

SENECIO [*Jacobæa*] corollis radiantibus, foliis pinnato-lyratis, laciniis lacinulatis, caule erecto. *Hudsoni Fl.* 316.

Classis Linnæi Syngenefia: Polygamia Superflua.
Common Ragwort.

In meadows and pastures; also in places uncultivated, and by the road side: common.

It flowers in July.

IRIS palustris lutea. *Raii Syn.* 374.

IRIS [*Pseudacorus*] corollis imberbibus, petalis interioribus stigmatibus minoribus, foliis ensiformibus. *Hudsoni Fl.* 13.

Classis Linnæi Triandria Monogynia.

Yellow Water Flower-de-luce.

In moist boggy places, and in rivers: common. Found in the river Rhodon, near Woodford-Bridge, and in other parts of it, plentifully.

It flowers in June.

IRIS sylvestris quam Xyrim vocant. *Raii Syn.* 375.

IRIS [*fætidissima*] corollis imberbibus, petalis interioribus patentissimis, caule unangulato, foliis ensiformibus. *Hudsoni Fl.* 13.

Classis Linnæi Triandria Monogynia.

Stinking Gladdon, or Gladwyn.

Under hedges, and on the sides of ditches. Found under a hedge in the lane leading from Chingford Hatch, to Mr. Lloyd's, Friday Hill House; and near Mr. Warner's Gravel-pit Pond: not common.

It flowers in June or July.

JUNIPERUS vulgaris baccis parvis purpureis.

Raii Syn. 444.

JUNIPERUS [*communis*] foliis ternis patentibus mucronatis, bacca longioribus. *Hudsoni Fl.*

372.

Classis Linnæi Dioecia Monodelphia.

The common Juniper.

On heaths and commons in dry, chalky or sandy places: common. In this part of Essex, uncommon; but found on the Forest between Waltham Abbey and Epping.

It flowers in May.

K.

KNAWEL. *Raii Syn.* 159.

SCLERANTHUS [*annuus*] calycibus fructus patulis. *Hudsoni Fl.* 160.

Classis Linnæi Decandria Digynia.

German Knotgrafs, or annual Knavel.

In barren, dry, sandy fields : very common.

It flowers in August.

L.

LACTUCA aguina seu Valerianella foliis ferratis.

Raii Syn. 201.

VALERIANA [*Locusta*] floribus triandris, caule dichotomo, foliis linearibus. *Hudsoni Fl.*

12.

Classis Linnæi Triandria Monogynia.

Small Corn Sallet, or Valerian with jagged leaves; *by some*, Small Lamb's-lettuce.

In fields, and among corn: not uncommon.

It flowers in April or May.

LACTUCA sylvestris major, odore Opii. *Raii Syn.* 161.

LACTUCA [*virosa*] foliis verticalibus carina aculeatis. *Hudsoni Fl.* 295.

Classis Linnæi Syngenesia: Polygamia Æqualis.

Greater strong scented wild Lettuce.

On banks, under hedges, and by the sides of fields.

It flowers in July or August.

LAMIUM album. *Raii Syn.* 240.

LAMIUM [*album*] foliis cordatis acuminatis serratis petiolatis. *Hudsoni Fl.* 225.

Classis Linnæi Didynamia Gymnospermia.

White Archangel, or dead Nettle.

Under hedges in uncultivated places, and by the sides of fields : very common.

It flowers in May.

LAMIUM cannabinum floribus albis, verticillis purpurascens. *Raii Syn.* 241.

GALEOPSIS [*Tetrahit*] internodiis caulinis superne incrassatis, verticillis summis subcontiguis. *Hudsoni Fl.* 226. β .

Classis Linnæi Didynamia Gymnospermia.

White flowered hemp-leaved dead Nettle.

In fields and among corn : common.

It flowers in August.

LAMIUM cannabinum flore amplo luteo, labio purpureo. *Raii Syn.* 241.

GALEOPSIS [*Tetrahit*] internodiis caulinis superne incrassatis, verticillis summis subcontiguis. *Hudsoni Fl.* 226.

Classis Linnæi Didynamia Gymnospermia.

Fair flowered Nettle-hemp; or rather, Hemp-leaved dead Nettle, with a parti-coloured flower.

In corn fields. Found in a corn field, at the bottom of Ribton Lane : not common.

It flowers in August.

LAMIUM cannabino folio vulgare. *Raii Syn.* 240.

GALEOPSIS [*Tetrahit*] internodiis caulinis superne incrassatis, verticillis summis subcontiguis. *Hudsoni Fl.* 226.

Classis Linnæi Didynamia Gymnospermia.

Nettle-hemp, or Hemp leaved dead Nettle.

In fields and among corn: not uncommon.

It flowers in August.

LAMIUM folio caulem ambiente majus et minus. *Raii Syn.* 240.

LAMIUM [*amplexicaule*] foliis floribus sessilibus amplexicaulibus obtusis. *Hudsoni Fl.* 225.

Classis Linnæi Didynamia Gymnospermia.

Great Henbit.

In corn fields, and often in cultivated ground and gardens: not uncommon.

It flowers in June.

LAMIUM luteum. *Raii Syn.* 242.

GALEOPSIS [*Galeobdolan*] verticillis sexfloris: involucro tetraphyllo. *Hudsoni Fl.* 226.

Classis Linnæi Didynamia Gymnospermia.

Yellow Archangel, dead Nettle, or yellow Nettle-hemp.

In woods, under hedges, and in shady places: common.

It flowers in May.

WOODFORDIENSES. 79

LAMIUM rubrum. *Raii Syn.* 140.

LAMIUM [*rubrum*] foliis cordatis, obtusis petiolatis. *Hudsoni Fl.* 225.

Classis Linnæi Didynamia Gymnospermia.

Red Archangel or dead Nettle.

On rubbish, and on the sides of fields : common.

It flowers in May.

LAMPSANA. *Raii Syn.* 173.

LAMPSANA [*communis*] calycibus fructus angulatis, pedunculis tenuibus ramosissimis. *Hudsoni Fl.* 303.

Classis Linnæi Syngenesia : Polygyamia Æqualis.

Nipplewort.

Under hedges, on rubbish, and in cultivated ground : not uncommon.

It flowers in June and July.

LAPATHUM maximum aquaticum, sive Hydro-lapathum: *Raii Syn.* 140.

RUMEX [*Britannica*] floribus hermaphroditis : valvulis integerrimis, omnibus graniferis, foliis lanceolatis vaginis obsoletis. *Hudsoni Fl.* 135.

Classis Linnæi Hexandria Trigynia.

Great Water Dock.

In ditches and boggy places : very common.

Found in different parts of the River Rhodon.

It flowers in July and August.

LAPATHUM acetosum repens lanceolatum. *Raii Syn.* 143.

RUMEX [*acetosella*] floribus dioicis, foliis lanceolato-hastatis. *Hudsoni Fl.* 136.

Classis Linnæi Hexandria Trigynia.
Sheeps-Sorrel.

On dry, sandy, ground, and on gravel: very common.

It flowers in May, June, or July.

LAPATHUM acetosum vulgare. *Raii Syn.* 143.

RUMEX [*acetosa*] floribus dioicis, foliis oblongis sagittatis. *Hudsoni Fl.* 136.

Classis Linnæi Hexandria Trigynia.
Common Sorrel.

In meadows, and pastures: common.

It flowers in May or June.

LAPATHUM acutum. *Raii Syn.* 142.

RUMEX [*acutus*] floribus hermaphroditis; valvulis dentatis graniferis, foliis cordato oblongis. *Hudsoni Fl.* 134.

Classis Linnæi Hexandria Trigynia.
Sharp-pointed Dock.

In uncultivated places, woods, and on swamps: very common.

It flowers in June.

WOODFORDIENSES. 81

LAPATHUM acutum minimum. *Raii Syn.* 141.

AN RUMICIS [*acuti*] *Hudsoni Fl.* 134. varietas.

Classis Linnæi Hexandria Trigynia.

The lesser sharp pointed Dock.

In woods, and moist grounds : common.

It flowers in June.

LAPATHUM folio acuto crispo. *Raii Syn.* 141.

RUMEX [*crispus*] floribus hermaphroditis : valvulis integris graniferis, foliis lanceolatis undulatis acutis. *Hudsoni Fl.* 134.

Classis Linnæi Hexandria Trigynia.

Sharp pointed Dock, with curled leaves.

In dry fields, and by the road side.

It flowers in June or July.

LAPATHUM viride. *Raii Syn.* 141.

AN RUMICIS [*acuti*] *Hudsoni Fl.* 134. varietas.

Classis Linnæi Hexandria Trigynia.

The greener leaved Dock.

In shady woods, and moist ground : common.

It flowers in June.

LAPATHUM vulgare folio obtuso. *Raii Syn.*

141.

RUMEX [*obtusifolium*] floribus hermaphroditis; valvulis dentatis, foliis cordato-oblongis, obtusiunculis crenulatis. *Hudsoni Fl.* 134.

Classis Linnæi Hexandria Trigynia.

Common broad-leaved Dock.

In moist places, and by the road side: very common.

It flowers in July or August.

LAPPA major Arctium Dioscoridis. *Raii Syn.*

197.

ARCTIUM [*Lappa*] foliis cordatis inermibus petiolatis. *Hudsoni Fl.* 304. δ .

Classis Linnæi Syngenesia: Polygamia Æqualis.

Burdock or Clot-burr.

Under hedges, and by the road side, every where: very common.

It flowers in July or August.

LAPPA major capitulo glabro maximo. *Raii*

Syn. 196.

ARCTIUM [*Lappa*] foliis cordatis inermibus petiolatis. *Hudsoni Fl.* 304. β .

Classis Linnæi Syngenesia: Polygamia Æqualis.

Taller Burdock, or Clot-burr, with a purple flower.

Under hedges, and by the road side: not uncommon.

It flowers in July or August.

LAPPA major capitulis parvis glabris. *Raii Syn.* 197.

ARCTIUM [*Lappa*] foliis cordatis inermibus petiolatis. *Hudsoni Fl.* 304. γ.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Burdock, or Clot-burr, with smaller heads.

Under hedges, and by the road side. Found in the lane leading to Muncombe.

It flowers in July or August.

LAPPA major montana, capitulis tomentosis, five
Arctium Dioscoridis. *Raii Syn.* 197.

ARCTIUM [*Lappa*] foliis cordatis inermibus petiolatis. *Hudsoni Fl.* 304. δ.

Classis Linnæi Syngenesia: Polygamia Æqualis.
Taller Burdock, or Clot-burr, with woolly heads.

By the road side: not uncommon.

It flowers in July or August.

LATHYRUS lutens sylvestris dumetorum. *Raii Syn.* 320.

LATHYRUS [*pratensis*] pedunculis multifloris, cirrhis diphyllis: foliis lanceolatis: cirrhis simplicissimis. *Hudsoni Fl.* 277.

Classis Linnæi Diadelphia Decandria.

Tare everlasting, or common yellow bastard Vetchling.

In woods, meadows, pastures, and on hedges: common.

It flowers in July or August.

LAUREOLA. *Raii Syn.* 465.

DAPHNE [*Laureola*] racemis axillaribus, foliis lanceolatis glabris. *Hudsoni Fl.* 145.

Classis Linnæi Octandria Monogynia.

Dwarf Laurel, or Spurge Laurel.

In woods and under hedges; in most places on the Forest: not uncommon.

It flowers in March or April.

LENS palustris. *Raii Syn.* 129.

LEMNA [*minor*] foliis sessilibus utrinque planiusculis, radicibus solitariis. *Hudsoni Fl.* 345.

Classis Linnæi Monoecia Diandria.

Ducks-meat.

In ponds and standing waters: very common.

It flowers in June.

LENTICULA aquatica trifulca. *Raii Syn.* 129.

LEMNA [*trifulca*] foliis petiolatis lanceolatis, *Hudsoni Fl.* 344.

Classis Linnæi Monoecia Triandria.

Ivy leaved Duck's-meat.

In ponds, ditches, and stagnated waters: not uncommon.

It flowers in May or June.

WOODFORDIENSES. 85

LENTICULA palustris major. *Raii Syn.* 129.

LEMNA [*polyrrhiza*] foliis sessilibus, radicibus confertis. *Hudsoni Fl.* 345.

Classis Linnæi Monoecia Diandria.

The greater Ducks-meat.

In ponds, ditches, and stagnated waters: common.

It flowers in June or July.

LEUCANTHEMUM vulgare. *Raii Syn.* 184.

CHRYSANTHEMUM [*Leucanthemum*] foliis amplexicaulibus oblongis: superne serratis: inferne dentatis. *Hudsoni Fl.* 321.

Classis Linnæi Syngenesia: Polygamia Superflua.

The greater Daisie, or Ox Eye.

In meadows, dry pastures, and among corn: not uncommon.

It flowers in May.

LEUCOJUM luteum, vulgo Cheiri, flore simplici.

Raii Syn. 291.

CHEIRANTHUS [*Cheiri*] foliis lanceolatis acutis glabris ramis angulatis. *Hudsoni Fl.* 250.

Classis Linnæi Tetradynamia Siliquosa.

Wall Flower, or Wild Cheir.

On walls, and on the roofs of houses; also, on rubbish: not uncommon.

It flowers in April or May.

LIGUSTRUM *Raii* Syn. 465.

LIGUSTRUM [*vulgare*] *Hudsoni* Fl. 3.

Classis Linnæi Diandrya Monogynia.

Privet, or Prim.

In woods and hedges : not uncommon.

It flowers in May.

LILIUM convallium. *Raii* Syn. 264.

CONVALLARIA [*majalis*] scapo nudo. *Hudsoni* Fl. 126.

Classis Linnæi Hexandria Monogynia.

Lily-convally, or May Lily.

On the shady parts of heaths, or in woods ; not common in these parts : but found opposite High Beech, near The King's Oak, in plenty.

It flowers in May.

LINARIA *Elatine dicta folio acuminato.* *Raii* Syn. * 282.

ANTIRRHINUM [*Elatine*] foliis hastatis alternis, caulibus procumbentibus, corollis calcaratis. *Hudsoni* Fl. 237.

Classis Linnæi Didynamia Angiospermia. Sharp-pointed Fluellin.

In corn fields : not uncommon.

It flowers in August or September.

LINARIA hederaceo folio glabro, seu Cymbalaria vulgaris. *Raii Syn.* * 282.

ANTIRRHINUM [*Cymbalaria*] foliis cordatis quinquelobis alternis, caulibus procumbentibus. *Hudsoni Fl.* 237.

Classis Linnæi Didynamia Angiospermia.

Ivy-leaved Toad-flax.

On walls and houses : not common.

It flowers from June to October.

LINARIA lutea vulgaris. *Raii Syn.* * 281.

ANTIRRHINUM [*Linaria*] foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. *Hudsoni Fl.* 238.

Classis Linnæi Didynamia Angiospermia.

Common yellow Toad-flax.

In barren meadows, on the sides of fields, and by the road side on banks : common.

It flowers in July or August.

LINUM sylvestre catharticum. *Raii Syn.* 362.

LINUM [*catharticum*] foliis oppositis ovato lanceolatis, caule dichotomo, corollis acutis. *Hudsoni Fl.* 116.

Classis Linnæi Pentandria Pentagynia.

Purging or wild Dwarf Flax. Mill-mountain.

In dry fields, and on banks. Found on a bank near Muncombe : not very common.

It flowers in May or June.

LONCHITIS aspera. *Raii Syn.* 118.

OSMUNDA [*Spicant*] frondibus lanceolatis pinatifidis: laciniis conniventibus integerrimis parallelis. *Hudsoni Fl.* 382.

Classis Linnæi Cryptogamia Filices.

Rough Spleenwort.

In moist woody places, and on banks, under hedges. Found on the side of a little running stream between Mr. Norris's, opposite The Eight Mile Stone, and Woodford Bridge: not very common.

It flowers in July.

LOTI corniculatæ major species. *Raii Syn.* 334.

LOTUS [*corniculata*] capitulis depressis caulibus decumbentibus, leguminibus cylindricis patentibus. *Hudsoni Fl.* 228. γ.

Classis Linnæi Diadelphia Decandria.

The greater Bird's Foot Trefoil.

In moist woods, meadows, and under hedges: not uncommon.

It flowers from June to August.

LOTUS corniculata glabra minor. *Raii Syn.* 334.

LOTUS [*corniculata*] capitulis depressis, caulibus decumbentibus, leguminibus cylindricis patentibus. *Hudsoni Fl.* 288.

Classis Linnæi Diadelphia Decandria.

Bird's Foot Trefoil.

In pastures and dry fields; also on the Forest: very common.

It flowers in June or July.

LUPULUS mas et fœmina. *Raii Syn.* 137.

HUMULUS [*Lupulus*] *Hudsoni Fl.* 369.

Classis Linnæi Dioecia Pentandria.

Hops, the male and female.

In hedges, in moist places : not uncommon.

It flowers in July.

LUTEOLA. *Raii Syn.* 366.

RESEDA [*Luteola*] foliis lanceolatis integris,
calycibus quadrifidis. *Hudsoni Fl.* 181.

Classis Linnæi Dodecandria Trigynia.

Wild Woad, yellow Weed, or Dyers Weed.

*In barren uncultivated places, and on walls :
not uncommon : sometimes in fields. Found on the
side of the hedge, the bottom of Sakes, commonly
called Snake's Lane, near Woodford Bridge,
and on a dry bank almost opposite the five mile
stone, Laytonstone.*

It flowers in June.

LYCHNIS sylvestris, quæ Behn album vulgo.
Raii Syn. 337.

CUCUBULUS [*Behen*] calycibus subglobosis gla-
bris reticulato-venosis capsulis trilocularibus,
corollis subnudis. *Hudsoni Fl.* 163.

Classis Linnæi Decandria Trigynia.

Spatling Popy, White Bottle, Bladder-Campion,
or white Corn-Campion.

*In dry pasture land, and in corn fields : not
uncommon.*

It flowers in July.

LYCHNIS plumaria sylvestris simplex. *Raii Syn.* 338.

LYCHNIS [*Flos cuculi*] petalis quadrifidis, fructu subrotundo. *Hudsoni Fl.* 174.

Classis Linnæi Decandria Pentagynia.

Meadow Pink, Wild Williams, Cuckow Flower.

In meadows and moist pastures : common.

It flowers in June.

LYCHNIS Saponaria dicta. *Raii Syn.* 339.

SAPONARIA [*officinalis*] calycibus cylindricis foliis ovato lanceolatis. *Hudsoni Fl.* 160.

Classis Linnæi Decandria Digynia.

Common Soap wort.

Under hedges, and walls of gardens ; but not common. Found under the wall of the garden of Friday Hill House, Mr. Lloyd's, and in Low-Layton Church-yard.

It flowers in July or August.

LYCHNIS segetum-major. *Raii Syn.* 338.

AGROSTEMMA [*Githaco*] hirsuta, calycibus corollam æquantibus, petalis integris nudis. *Hudsoni Fl.* 173.

Classis Linnæi Decandria Pentagynia.

Cockle.

In corn fields : common.

It flowers in June.

WOODFORDIENSES. 91

LYCHNIS sylvestris albo flore. *Raii Syn.* 339.

LYCHNIS [*dioica*] floribus dioicis. *Hudsoni Fl.*

174.

Classis Linnæi Decandria Pentagynia.

Wild white Campion.

In woods, under hedges, and on the sides of fields : not very uncommon ; but not found so often, as that with a red flower.

It flowers in May, June, and July.

LYCHNIS sylvestris flore rubello. *Raii Syn.* 339.

LYCHNIS [*dioica*] floribus dioicis. *Hudsoni Fl.*

174.

Classis Linnæi Decandria Pentagynia. β.

Red flowered wild Campion ; *when double and in gardens, called Batchelor's Buttons.*

In woods, under hedges, and on the sides of fields : very common.

It flowers in May, June, and July.

LYCOPUS palustris glaber. *Raii Syn.* 236.

LYCOPUS [*Europæus*] foliis sinuato-ferratis.

Hudsoni Fl. 8.

Classis Linnæi Diandria Monogynia.

Water Horehound.

On the sides of rivers, and on moist, boggy places : common. Found in plenty on the side of the foot-path, between Woodford Row and Woodford Church.

It flowers in July.

LYSIMACHIA lutea. *Raii Syn.* 282.

LYSIMACHIA [*vulgaris*] paniculata, racemis terminalibus. *Hudsoni Fl.* 72.

Classis Linnæi Pentandria Monogynia.

Yellow Willow-herb, or Loose-strife.

In watery places, and on the banks of rivers : but in general, not very common. Found on the banks of the river Rhodon near Luxborough-House, Sir Edward Walpole's,

It flowers in July or August.

LYSIMACHIA campestris. *Raii Syn.* 311.

EPILOBIUM [*montanum*] foliis oppositis ovatis dentatis. *Hudsoni Fl.* 141.

Classis Linnæi Octandria Monogynia.

The greater smooth leaved Willow-herb, or Loose-strife.

By ditch sides, under hedges ; and in the moist woody parts of the Forest : common.

It flowers in June or July.

LYSIMACHIA filiquosa glabra media sive minor. *Raii Syn.* 311.

EPILOBIUM [*tetragonum*] foliis lanceolatis-linearibus denticulatis oppositis, caule tetragono. *Hudsoni Fl.* 141.

Classis Linnæi Octandria Monogynia.

Middle smooth leaved Willow-herb, or Loose-strife.

In moist ditches, and near rivers : common.

It flowers in July.

WOODFORDIENSES. 93

LYSIMACHIA filiquosa glabra minor angustifolia. *Raii Syn.* 311.

EPILOBIUM [*palustre*] oppositis foliis lanceolatis integerrimis, petalis emarginatis caule erecto. *Hudsoni Fl.* 141.

Classis Linnæi Octandria Monogynia.

The least smooth coddled Loofestripe, or Willow-herb.

In marshy places : common.

It flowers in July.

LYSIMACHIA filiquosa hirsuta magno flore.
Raii Syn. 311.

EPILOBIUM [*ramosum*] foliis lanceolatis ferratis subdecurrentibus inferioribus oppositis, caule ramoso. *Hudsoni Fl.* 141.

Classis Linnæi Octandria Monogynia.

Great flowered Willow-herb, called Codlings and Cream.

In moist places under hedges, and by the sides of rivers. Found on the sides of the river Rhodon in plenty.

It flowers in July.

LYSIMACHIA filiquosa hirsuta parvo flore. *Raii Syn.* 311.

EPILOBIUM [*hirsutum*] foliis lanceolatis undulato-ferratis decurrentibus : inferioribus oppositis, caule sub simplici racemo terminali. *Hudsoni Fl.* 140.

Classis Linnæi Octandria Monogynia.

The lesser hairy coddled Loofestriſe, or Willow-herb, with small flowers.

In moist places; and, on the sides of rivers : common.

It flowers in July.

LYSIMACHIA speciosa, quibusdam Onagra dicta. *Raii Syn.* 310.

EPILOBIUM [*angustifolium*] foliis sparsis linearilanceolatis. *Hudsoni Fl.* 140.

Classis Linnæi Octandria Monogynia.

Rosebay Willow-herb.

In woods and under hedges; sometimes in meadows; but in these parts, uncommon. Found in a meadow, near the river Rhodon, not far from Luxborough House, Sir Edward Walpole's.

It flowers in July and August.

M.

MALVA sylvestris minor. *Raii Syn.* 251.

MALVA [*rotundifolia*] caule prostrato, foliis cordato-orbiculatis obsolete quinquelobis.

Hudsoni Fl. 268.

Classis Linnæi Monadelphia Polyandria.

Small wild Mallow, or Dwarf Mallow.

On dry banks by the road side: common.

It flowers from July to October.

MALVA vulgaris. *Raii Syn.* 251.

MALVA [*sylvestris*] caule erecto herbaceo, foliis lobatis obtusis, pedunculis petiolisque pilosis. *Hudsoni Fl.* 268.

Classis Linnæi Monadelphia Polyandria.

Common Mallow.

Under hedges, by the way-side, and on rubbish: very common.

It flowers from June to October.

MALUS

MALUS sylvestris. *Raii Syn.* 452.

PYRUS [*malus*] foliis ferratis umbellis sessilibus.
Hudsoni Fl. 189.

Classis Linnæi Icosandria Pentagynia.

The Crab Tree or Wilding.

In woods and hedges. Found on the Forest, opposite Hagger Lane, near the Gibbets, and in a Hedge by the river Rhodon, near Luxborough House, Sir Edward Walpole's: not uncommon.

It flowers in May.

MATRICARIA. *Raii Syn.* 187.

MATRICARIA [*Parthenium*] foliis compositis planis; foliolis ovatis incisiss, pedunculis ramosis. *Hudsoni Fl.* 321.

Classis Linnæi Syngenesia: Polygamia Superflua.
Fever-few.

Under hedges, among rubbish, and sometimes upon old walls: common.

It flowers in June.

MELAMPYRUM sylvaticum flore luteo, five Satureia lutea sylvestris. *Raii Syn.* * 286.

MELAMPYRUM [*sylvaticum*] corollis hiantibus.
Hudsoni Fl. 236.

Classis Linnæi Didynamia Angiospermia.

Common wild Cow Wheat, with a yellow flower.

In woods, and among bushes: very common.

It flowers in June, July, or August.

MELILOTUS vulgaris. *Raii Syn.* 331.

TRIFOLIUM [*melilotus officinalis*] leguminibus racemosis nudis dispermis, caule erecto. *Hudsoni Fl.* 282.

Classis Linnæi Diadelphia Decandria.

Common Melilot.

In fields and under hedges : among corn : common.

It flowers in July or August.

MENIANTHES palustre triphyllum latifolium et angustifolium. *Raii Syn.* 285.

MENYANTHES [*trifolia*] foliis ternatis. *Hudsoni Fl.* 71.

Classis Linnæi Pentandria Monogynia.

Marsh Trefoil or Buck-bean.

In marshes and moist meadows : common.

It flowers in June or July.

MENTHA aquatica, seu Sisymbrium. *Raii Syn.* 233.

MENTHA [*aquatica*] floribus capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus. *Hudsoni Fl.* 223.

Classis Linnæi Didynamia Gymnospermia.

Water Mint.

In moist places, and on the sides of rivers : very common.

It flowers in July and August.

MENTHA seu Calamintha aquatica. *Raii Syn.*
232.

MENTHA [*arvensis*] floribus verticillatis foliis
ovatis acutis serratis staminibus corollam
æquantibus. *Hudsoni Fl.* 223.

Classis Linnæi Didynamia Gymnospermia.

Water Mint with whorled Coronets.

In moist meadows, and in places where stagnated waters have remained through the winter, and have been dried up in summer: not uncommon.

It flowers in August and September.

MENTHA spicis brevioribus, et habitioribus,
foliis Menthæ fuscæ, sapore fervido piperis.
Raii Syn. 234.

MENTHA [*piperita*] floribus capitatis solitariis
terminalibus, foliis ovato-oblongis, serratis,
staminibus corolla brevioribus. *Hudsoni Fl.*
222.

Classis Linnæi Didynamia Gymnospermia.

Pepper-Mint.

In moist places, usually near rivers. Found in a lane which leads from Dr. Wilkinson's House in Woodford, to The Wind Mill: very uncommon.

It flowers in August.

WOODFORDIENSES. 99

MERCURIALIS annua glabra vulgaris. Mercurialis mas et fœmina. *Raii Syn.* 139.

MERCURIALIS [*annua*] caule brachiato, foliis glabris. *Hudsoni Fl.* 371.

Classis Linnæi Dioecia Enneandria.

French Mercury, the male and female.

On rubbish, and often on cultivated land: not uncommon.

It flowers in September.

MERCURIALIS perennis repens, Cynocrambe dicta. *Raii Syn.* 138.

MERCURIALIS [*perennis*] caule simplicissimo, foliis scabris. *Hudsoni Fl.* 371.

Classis Linnæi Dioecia Enneandria.

Dogs Mercury.

In woods and under hedges: common.

It flowers in April and May.

MESPILUS apii folio sylvestris non spinosa, seu Sorbus terminalis. *Raii Syn.* 453.

CRATÆGUS [*torminalis*] foliis cordatis septangulis: lobis infimis divaricatis. *Hudsoni Fl.* 188.

Classis Linnæi Icosandria Digynia.

The common wild Service Tree, or Sorb.

In woods and hedges. Found in Sake's, commonly called Snake's Lane. On a bank, between Mr. Warner's Fields and Ribton Lane, and in a wood, called The Sale, in the parish of Walthamstow.

It flowers in April or May.

MESPILUS apii folio fylvestris spinosa, five Oxycantha. *Raii Syn.* 453.

CRATÆGUS [*Oxycantha*] foliis obtusis subtrifidis ferratis. *Hudsoni Fl.* 188.

Classis Linnæi Icosandria Digynia.

The White Thorn, or Hawthorn. Often called Quick.

In hedges: very common every where: sometimes in Woods, on heaths and commons. Found without thorns, and with a large leaf, in a hedge on the side of the river Rhodon, between Wray House and Luxborough House, opposite the path that leads to Muncombe.

It flowers in May.

MILLEFOLIUM vulgare. *Raii Syn.* 183.

ACHILLEA [*millefolium*] foliis bipinnatis nudis: laciniis linearibus dentatis. *Hudsoni Fl.* 324.

Classis Linnæi Syngenesia: Polygamia Superflua.

Common Yarrow, Milfoil, or Nose-bleed.

In meadows and pasture land: very common. Found also sometimes with a purple flower; though it has usually a white one.

It flowers in May, June, or July.

MOLLUGO montana minor Gallio albo similis.

Raii Syn. 224.

GALLIUM [*montanum*] foliis fenis obovatis, mucronatis glabris, caule prostrato, pedunculis ternatis. *Hudsoni Fl.* 56.

Classis Linnæi Tetrandria Monogynia.

Small Mountain Bastard Madder. Mountain Ladies Bedstraw.

On dry ground, and hilly situations: not uncommon.

It flowers in July.

MOLLUGO vulgator. *Raii Syn.* 223.

GALLIUM [*mollugo*] foliis octonis, ovato-linearibus subserratis patentissimis mucronatis, caule flaccido, ramis patentibus. *Hudsoni Fl.* 56.

Classis Linnæi Tetrandria Monogynia.

Wild Madder, or great Bastard Madder.

On hedges, and in bushy places: very common.

It flowers in June or July.

MOLLUGINIS vulgatoris varietas minor. *Raii Syn.* 224.

GALLIUM [*palustre*] foliis quaternis obovatis inæqualibus, caulibus diffusis. *Hudsoni Fl.* 57.

Classis Linnæi Tetrandria Monogynia.

White Ladies Bedstraw.

On the sides of rivers, and in moist meadows: very common.

It flowers in July.

MYOSOTIS scorpioides hirsuta. *Raii Syn.* 229.

MYOSOTIS [*scorpioides*] feminibus nudis; foliorum apicibus callofis. *Hudsoni Fl.* 65.

Classis Linnæi Pentandria Monogynia.

Mouſe-ear. Scorpion-graſs.

In dry paſtures; ſometimes in corn fields, and in gardens, ſpontaneouſly: common.

It flowers from April to Auguſt.

MYOSOTIS scorpioides hirta minor. *Raii Syn.* 229.

MYOSOTIS [*scorpioides*] feminibus nudis; foliorum apicibus callofis. *Hudsoni Fl.* 65. 7.

Classis Linnæi Pentandria Monogynia.

Small yellow flowered Scorpion Graſs.

On walls, barren places, and on gravelly ſoils: not uncommon.

It flowers in April or May.

MYOSOTIS scorpioides palustris. *Raii Syn.* 229.

MYOSOTIS [*scorpioides*] feminibus nudis; foliorum apicibus callofis. *Hudsoni Fl.* 65. 8.

Classis Linnæi Pentandria Monogynia.

Mouſe-ear. Scorpion-graſs.

In barren, dry fields, und ſometimes on cultivated ground: not uncommon.

It flowers from April to Auguſt.

WOODFORDIENSES. 103

MYOSURUS. *Raii* Syn. 251.

MYOSURUS [*minimus*] foliis integris. *Hudsoni*
Fl. 118.

Classis Linnæi Pentandria Polygynia.

Moufe-tail.

In fields, on a gravelly soil: Found in the road between Woodford and Chingford Hatch: uncommon.

It flowers in April or May.

MYRRHIS fylvestris feminibus asperis. *Raii*
Syn. 220.

SCANDIX [*Anthriscus*] feminibus ovatis hispidis,
corollis uniformibus, caule lævi. *Hudsoni*
Fl. 108.

Classis Linnæi Pentandria Digynia.

Small Hemlock Chervil, with rough seeds.

Under hedges: common.

It flowers in May or June.

N

NAPUS sylvestris. *Raii Syn.* 295.

BRASSICA [*Napus*] radice caulescenti fusiformi. *Hudsoni Fl.* 253.

Classis Linnæi Tetradynamia Siliquosa.

Wild Navew.

Among corn, and on the sides of ditches.

It flowers in May.

NARCISSUS sylvestris pallidus calyce luteo. *Raii Syn.* 371.

NARCISSUS [*Pseudo-Narcissus*] spatha uniflora, nectarii limbo campanulato erecto petalis æquali. *Hudsoni Fl.* 123.

Classis Linnæi Hexandria Monogynia.

Wild English Daffodil.

On the sides of fields, chiefly in a moist soil: in woods, bushy places, and under plantations of trees. Found in the upper part of Mr. Warner's large field in plenty.

It flowers in March or April.

NASTURTIUM fupinum capfulis verrucofis : five
Coronopus Ruellii. *Raii Syn.* 304.

COCHLEARIA [*Coronopus*] foliis pinnatifidis.
Hudfoni Fl. 248.

Classis Linnæi Tetradynamia Siliculofa.
Swine's Creffes.

In moist places; on the fides of roads and paths. Found near a pond almost opposite the nine mile stone Woodford Row : and on the fide of the road to Woodford Bridge, near the eight mile stone : not very common.

It flowers in June, July, or Auguft.

NEPETA major vulgaris. *Raii Syn.* 237.

NEPETA [*Cataria*] floribus fpicatis verticillatis
fubpedicellatis, foliis petiolatis cordatis den-
tato ferratis. *Hudfoni Fl.* 220.

Classis Linnæi Didynamia Gymnofpermia.

Nep, Nap, or Cat-mint.

In fields, and under hedges. Found near Abridge : not very common.

It flowers in July.

NUMMULARIA. *Raii Syn.* 283.

LYSIMACHIA [*Nummularia*] foliis ovatis acutis,
floribus folitariis, caule procumbente. *Hud-
foni Fl.* 73.

Classis Linnæi Pentandria Monogynia.

Money-wort, or Herb Two-pence.

*In fields, and moist places near ditches. Found in Mr. Warner's large field, in plenty : not un-
common.*

It flowers in June.

NUMMULARIA minor flore purpurascente.

Raii Syn. 283.

LYSIMACHIA [*tenella*] foliis ovatis acutiusculis,
pedunculis folio longioribus, caule repente.

Hudsoni Fl. 73.

Classis Linnæi Pentandria Monogynia.

Purple flowered Money-wort.

In moist places on Heaths. Found on the Forest, opposite The Bald Faced Stag, going towards Chingford: not common.

It flowers in July or August.

NYMPHÆA alba. *Raii Syn.* 368.

NYMPHÆA [*alba*] foliis cordatis integerrimis
calyce quadrifido. *Hudsoni Fl.* 206.

Classis Linnæi Polyandria Monogynia.

Water Lily, with a white flower.

In rivers and standing waters. Found in the river Rhodon, near Luxborough House, Sir Edward Walpole's, and near Pissingford Bridge: not very common.

It flowers in July or August.

NYMPHÆA lutea. *Raii Syn.* 368.

NYMPHÆA [*lutea*] calyce magno pentaphyllo.
Hudsoni Fl. 206.

Classis Linnæi Polyandria Monogynia.

Water Lily, with a yellow flower.

In rivers and large standing waters. Found in many parts of the river Rhodon, particularly between Woodford Bridge and Luxborough House, Sir Edward Walpole's; and in a standing water by the road side, between the Old Mill and the eight mile stone.

It flowers in July or August.

NYM-

WOODFORDIENSES. 107

NYMPHÆA lutea minor flore fimbriato. *Raii*
Syn. 368.

MENYANTHES [*Nymphyoides*] foliis cordatis
integerrimis, corollis ciliatis. *Hudsoni Fl.* 71.
Classis Linnæi Pentandria Monogynia.

The lesser yellow Water Lily, with a fringed
flower.

*In rivers and large ditches. Found in the
river Rhodon, almost opposite the eight mile stone
in the road to Woodford Bridge, near the bridge,
and between that and Luxborough House, Sir
Edward Walpole's: not common.*

It flowers in June or July.

O.

OENANTHE aquatica. *Raii Syn.* 210.

OENANTHE [*fistulosa*] stolonifera, foliis caulinis pinnatis filiformibus fistulosis. *Hudsoni Fl.* 104.

Classis Linnæi Pentandria Digynia.

Water Dropwort.

In ditches, moist places, and by the sides of rivers: not very common.

It flowers in July.

OENANTHE aquatica triflora, caulibus fistulosis. *Raii Syn.* 210.

OENANTHE [*fistulosa*] stolonifera, foliis caulinis pinnatis filiformibus fistulosis. *Hudsoni Fl.* 104. β.

Classis Linnæi Pentandria Digynia.

The lesser Water-Dropwort.

In ditches, moist places, and by the sides of rivers: not very common.

It flowers in July.

WOODFORDIENSES. 109

OENANTHE Cicutæ facie Lobelii. *Raii Syn.*
210.

OENANTHE [*crocata*] foliis omnibus multifidis
obtusis subæqualibus. *Hudsoni Fl.* 105.

Classis Linnæi Pentandria Digynia.

Hemlock Dropwort.

*On the sides of rivers. Found on the banks
of the river Rhodon, near Luxborough House,
Sir Edward Walpole's; not very common.*

It flowers in June.

ONOBRYCHIS seu Caput Gallinaceum. *Raii
Syn.* 327.

HEDYSARUM [*Onobrychis*] foliis pinnatis legu-
minibus monospermis aculeatis, corollarum
alis brevissimis, caule elongato. *Hudsoni Fl.*
281.

Classis Linnæi Diadelphia Decandria.

Medick-Vetchling, or Cock's-head; *falsely cal-
led* Saint Foin.

*In chalky or gravelly fields. Found in a field
near Muncombe: not common.*

It flowers in July.

OPHIOGLOSSUM. *Raii Syn.* 128.

OPHIOGLOSSUM [*vulgatum*] fronde ovata. *Hud-
soni Fl.* 382.

Classis Linnæi Cryptogamia: Filices.

Adder's Tongue.

*In meadows, pastures, and sometimes in moist
places upon heaths. Found in a Field belonging
to Mr. Norris, almost opposite the eight mile stone
in Woodford: not common.*

It flowers in May.

OPULUS Ruellii. *Raii Syn.* 460.

VIBURNUM [*Opulus*] foliis ovatis, petiolatis glandulosis. *Hudsoni Fl.* 112.

Classis Linnæi Pentandria Trigynia.

Water Elder; called also, Cherry-wood.

In moist places in the woody part of the Forest. Found also in a hedge by the sides of the river Rhodon, between Wray - House, Sir James Wright's, and Luxborough House, Sir Edward Walpole's.

It flowers in May or June.

ORCHIS alba bifolia minor calcari oblongo.

Raii Syn. 380.

ORCHIS [*bifolia*] bulbis indivisis nectarii labio lanceolato integerrimo: cornu longissimo, petalis patentibus. *Hudsoni Fl.* 333.

Classis Linnæi Gynandria Diandria.

The lesser Butterfly Orchis.

In moist meadows: not very common.

It flowers in May or June.

ORCHIS morio mas, folis maculatis. *Raii Syn.* 376.

ORCHIS [*mascula*] bulbis indivisis, nectarii labio quadrilobo crenulato: cornu obtuso, petalis dorsalibus reflexis. *Hudsoni Fl.* 333.

Classis Linnæi Gynandria Diandria.

Male Orchis, or Male Fool's Stones.

In fields and meadows. Found in a field leading from Friday Hill House, Mr. Lloyd's, to Fairmead Bottom, in plenty.

It flowers in May.

ORCHIS.

WOODFORDIENSES. 111

ORCHIS morio foemina. *Raii Syn.* 377.

ORCHIS [*morio*] bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso, petalis omnibus conniventibus. *Hudsoni Fl.* 333.

Classis Linnæi Gynandria Diandria.

Female Orchis, or Female Fool Stones.

In Fields, and moist meadows. Found as the preceding, in plenty.

It flowers in May or June.

ORCHIS palmata pratensis latifolia, longis calcaribus. *Raii Syn.* 380.

ORCHIS [*latifolia*] bulbis subpalmatis rectis, nectarii cornu conico: labio trilobo lateralibus reflexis, bracteis flore longioribus. *Hudsoni Fl.* 335.

Classis Linnæi Gynandria Diandria.

Male handed Orchis, or Male handed Satyrion Royal.

In moist meadows, and on boggy ground.

It flowers in May or June.

ORCHIS palmata speciore thyrso folio maculato, sive palmata pratensis maculata. *Raii Syn.* 381.

ORCHIS [*maculata*] bulbis palmatis patentibus, nectarii cornu germinibus brevioribus: labio plano, petalis dorsalibus erectis. *Hudsoni Fl.* 335.

Classis Linnæi Gynandria Diandria.

Female handed Orchis, or Female Satyrion Royal.

In woods and meadows, on a dry soil.

It flowers in June.

ORCHIS spiralis alba odorata sive Triorchis.
Raii Syn. 378.

OPHRYS [*spiralis*] bulbis aggregatis, oblongis,
 caule subfolioso, floribus secundis, nectarii
 labio indiviso. *Hudsoni Fl.* 338.

Classis Linnæi Gynandria Diandria.

Triple-Ladies Traces.

*In dry fields. Found on the Forest, near The
 Bald-faced Stag, on the right hand of the road,
 not far from the eight mile stone, nearer London :
 not common.*

It flowers in August or September.

ORIGANUM vulgare spontaneum. *Raii Syn.*
 236.

ORIGANUM [*vulgare*] spicis subrotundis pani-
 culatis conglomeratis, bracteis calyce longi-
 oribus ovatis. *Hudsoni Fl.* 229.

Classis Linnæi Didynamia Gymnospermia.

Wild Marjoram.

*In the fields, under hedges, and in bushy places :
 common.*

It flowers in July.

ORNITHOPODIUM radice nodosa. *Raii Syn.* 326.

ORNITHOPUS [*perpusillus*] foliis pinnatis, le-
 guminibus subarcuatis. *Hudsoni Fl.* 280.

Classis Linnæi Diadelphia Decandria.

Bird's Foot.

*In the fields, on sandy or gravelly soils. Found
 on the Forest near Mr. Moxon's House : common.*

It flowers from May to August.

WOODFORDIENSES. 113

OROBANCHE major Caryophyllum olens. *Raii*
Syn. * 288.

OROBANCHE [*major*] caule simplicissimo pubescente, staminibus subexsertis. *Hudsoni Fl.*
232.

Classis Linnæi Dydynamia Gymnospermia.
Broom-rape.

In meadows, by the road side, and on the Forest, in dry soils. Found near the Wind-mill, the Seven Mile Stone in the Woodford Bridge road, and in a field where Broom grows, on the left hand of Hagger-lane, leading to Waltham-stow, in great plenty.

It flowers in May or June.

OROBUS sylvaticus foliis oblongis glabris. *Raii*
Syn. 324.

OROBUS [*tuberosus*] foliis pinnatis lanceolatis stipulis semi-sagittatis caule simplici. *Hudsoni Fl.* 274.

Classis Linnæi Diadelphia Decandria.

Wood Pea or Heath Pea.

In woods and copses: common.

It flowers in May.

OSTRYA Ulmo similis, fructu in umbilicis foliaceis. *Raii Syn.* 451.

CARPINUS [*Betulus*] squamis strobilorum planis. *Hudsoni Fl.* 360.

Classis Linnæi Monoecia Polyandria.

The Horn, or Hard-beam Tree, called in some places, the Horse-beech or Horn-beech, from some likeness of the leaves to the Beech.

In all parts of the Forest; and in hedges; very common.

It flowers in May.

OXYS alba. *Raii Syn.* * 281.

OXALIS [*Acetosella*] scapo unifloro, foliis ternatis squamoso-articulatis. *Hudsoni Fl.* 173.

Classis Linnæi Decandria Pentagynia.

Wood Sorrel.

In woods, and in shady places on heaths. Found in the woody part of the Forest, between Woodford Row and Chingford Hatch: not uncommon.

It flowers in June.

P.

PAPAVER corniculatum luteum, Chelidonia dictum. *Raii Syn.* 309.

CHELIDONIUM [*majus*] pedunculis umbellatis. *Hudsoni Fl.* 201.

Classis Linnæi Polyandria Monogynia.

The greater Celandine.

In dry ground and among rubbish. Found on the wall before Mr. Godfrey's House in Woodford, and in a Lane by the side of Mr. Raymond's House in Chigwell: not very common.

It flowers in May or June.

PAPAVER laciniato folio, capitulo breviori glabro annum, Rhoëas dictum. *Raii Syn.* 308.

PAPAVER [*Rhoëas*] capsulis glabris globosis, caule piloso multifloro, foliis pinnatifidis incis. *Hudsoni Fl.* 203.

Classis Linnæi Polyandria Monogynia.

Red Poppy, Corn Poppy, round smooth headed Poppy, or Corn Rose.

In corn fields; and by the road side: common

It flowers in June or July,

PAPAYER laciniato folio, capitulo hispido longiore. *Raii Syn.* 308.

PAPAYER [*Argemone*] capsulis clavatis hispidis, caule folioso multifloro. *Hudsoni Fl.* 203.

Classis Linnæi Polyandria Monogynia.

Long rough headed bastard Poppy.

In corn fields : not uncommon.

It flowers in June.

PARIETARIA. *Raii Syn.* 158.

PARIETARIA [*officinalis*] foliis lanceolato ovatis, pedunculis dichotomis, calycibus diphyllis. *Hudsoni Fl.* 376.

Classis Linnæi Polygamia Monoecia.

Pellitory of the Wall.

On walls and rubbish. Found on the east wall of Woodford Church-yard in plenty, and on the wall before Mr. Columbier's House, in Chigwell : not uncommon.

It flowers from May to August.

PARONYCHIA vulgaris. *Raii Syn.* 292.

DRABA [*verna*] scapis nudis, foliis subferratis. *Hudsoni Fl.* 243.

Classis Linnæi Tetradynamia Siliculosa.

Common Whitlow-grass.

On the Forest, in dry ground ; and on the turf in lawns or gardens : common.

It flowers in April.

WOODFORDIENSES. 117

PASTINACA sylvestris latifolia. *Raii Syn.* 206.

PASTINACA [*sativa*] foliis simpliciter pinnatis.
Hudsoni Fl. 109.

Classis Linnæi Pentandria Digynia.

Wild Parsnep.

On the sides of fields: very uncommon

It flowers in July.

PEDICULARIS, seu *Christa galli lutea*. *Raii Syn.* * 284.

RHINANTHUS [*Christa galli*] corollâ labio superiore compresso brevior. *Hudsoni Fl.* 234.

Classis Linnæi Didynamia Angiospermia.

Yellow Rattle, or Cocks-comb.

In barren fields and meadows: common.

It flowers in June or July.

PEDICULARIS palustris rubra elatior. *Raii Syn.* * 284.

PEDICULARIS [*palustris*] caule ramoso, calycibus calloso-punctatis, corollis labio obliquis.
Hudsoni Fl. 236.

Classis Linnæi Didynamia Angiospermia.

Great Marsh Red Rattled Coxcomb. *Louf-wort.*

In marshes or moist meadows: not uncommon.

It flowers in June.

PEDICULARIS pratensis rubra vulgaris. *Raii Syn.* * 284.

PEDICULARIS [*sylvatica*] caule ramoso, calycibus oblongis angulatis lævibus, corollis labio cordato. *Hudsoni Fl.* 236.

Classis Linnæi Didynamia Angiospermia.

Red Rattle Coxcomb. Common Loufewort.

In fields, and on moist parts of the Forest: very common.

It flowers in May or June.

PENTAPHYLLUM vulgatissimum. *Raii Syn.* 225.

POTENTILLA [*reptans*] foliis digitatis, caule repente, pedunculis unifloris. *Hudsoni Fl.* 197.

Classis Linnæi Icosandria Polygynia.

Common Cinquefoil, or five-leaved Grass.

Found near paths, and on the road side: very common.

It flowers in June.

PENTAPHYLLUM erectum, foliis profundè sectis, subtus argenteis, flore luteo. *Raii Syn.* 255.

POTENTILLA [*argentea*] foliis quinatis cuneiformibus incisus subtus tomentosis, caule erecto. *Hudsoni Fl.* 196.

Classis Linnæi Icosandria Polygynia.

Tormentil Cinquefoil.

On sandy, barren soils: not very common.

It flowers in June.

WOODFORDIENSES. 119

PENTAPHYLLOIDES Argentina dicta. *Raii Syn.* 256.

POTENTILLA [*Argentina*] foliis pinnatis, serratis, caule repente. *Hudsoni Fl.* 195.

Classis Linnæi Icofandria Polygynia.

Wild Tansey or Silver Weed.

Near paths on the road side, and where stagnated waters have remained through the winter : very common.

It flowers in June or July.

PENTAPTEROPHYLLON aquaticum flosculis ad foliorum nodos. *Raii Syn.* 316.

MYRIOPHYLLUM [*verticillatum*] floribus omnibus verticillatis. *Hudsoni Fl.* 357.

Classis Linnæi Monoecia Polyandria.

Verticillated Water Millfoil.

In ditches and stagnated waters : common.

It flowers in July.

PERCEPIER Anglorum. *Raii Syn.* 159.

APHANES [*arvensis*] *Hudsoni Fl.* 60.

Classis Linnæi Tetrandria Digynia.

Pursley-Piert.

In dry fields, on gravelly soils, and among corn : common.

It flowers in May.

PERSICARIA folio subtus incano. *Raii Syn.*

145.

POLYGONUM [*Persicaria* γ] floribus hexandris digynis, spicis ovato oblongis, foliis lanceolatis, stipulis ciliatis. *Hudsoni Fl.* 147.

Classis Linnæi Octandria Digynia.

Dead or spotted Arsmart, with leaves hoary on the back side.

In meadows, ditches, and on moist soils; in corn fields: very common.

It flowers in August or September.

PERSICARIA latifolia geniculata, caulibus maculatis. *Raii Syn.* 145.

POLYGONUM [*Persicaria* ϵ] floribus hexandris digynis, spicis ovato oblongis, foliis lanceolatis, stipulis ciliatis. *Hudsoni Fl.* 147.

Classis Linnæi Octandria Trigynia.

Dead or spotted Arsmart, with spotted stalks.

In fields, and on rubbish: common.

It flowers in August or September.

PERSICARIA maculosa. *Raii Syn.* 145.

POLYGONUM [*Persicaria*] floribus hexandris digynis spicis ovato-oblongis, foliis lanceolatis, stipulis ciliatis. *Hudsoni Fl.* 147.

Classis Linnæi Octandria Trigynia.

Dead or Spotted Arsmart.

In meadows, fields, and ditches: very common.

It flowers in August or September.

PERSICARIA mitis major, foliis pallidioribus
D. Bobarti. *Raii Syn.* 145.

POLYGONUM [*pensylvanicum*] floribus octandris
digynis, pedunculis hispidis, foliis lanceolatis,
stipulis muticis. *Hudsoni Fl.* 148.

Classis Linnæi Octandria Trigynia.

Greater dead Arsmart, with pale leaves.

In fields, on dunghills, and rich soils: not very common.

It flowers in August.

PERSICARIA falicis folio perennis, Potomogiton
angustifolium dicta. *Raii Syn.* 145.

POLYGONUM [*amphibium*] floribus pentandris
semidigynis, staminibus corolla longioribus.
Hudsoni Fl. 147.

Classis Linnæi Octandria Trigynia.

Perennial Willow-leaved Arsmart.

In fields, on rubbish, in moist places. Found on the banks of ditches and rivers, and often floating on their waters; sometimes in dry fields: not uncommon.

It flowers in June or July.

PERSICARIA vulgaris acris seu Hydropiper.
Raii Syn. 144.

POLYGONUM [*Hydropiper*] floribus hexandris
semidigynis, foliis lanceolatis, stipulis sub-
muticis. *Hudsoni Fl.* 148.

Classis Linnæi Octandria Trigynia.

Lakeweed. Arsmart. Water Pepper.

In moist places, and wet ditches: common.

It flowers in July or August.

PETASITES. *Raii Syn.* 179.

TUSSILAGO [*Petasites*] thyrso ovato flosculis omnibus hermaphroditis. *Hudsoni Fl.* 315.

Classis Linnæi Syngenesia: Polygamia Superflua.

Common Butter-bur. Pestilent Wert.

In moist ground, or near rivers. Found near the river Rhodon: not very uncommon.

It flowers in March or April.

PHELLANDRIUM vel Cicuta aquatica quorundam. *Raii Syn.* 215.

PHELLANDRIUM [*aquaticum*] foliorum ramificationibus divaricatis. *Hudsoni Fl.* 107.

Classis Linnæi Pentandria Digynia.

Water Hemlock.

In moist ditches and rivers. Found in the river Rhodon, and in ditches, near the eighth mile stone, in the Woodford Bridge road: not uncommon.

It flowers in July.

PHYLLITIS seu Lingua cervina vulgò. *Raii Syn.* 116.

ASPLENIUM [*Scolopendrium*] frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. *Hudsoni Fl.* 384.

Classis Linnæi Cryptogamia: Filices.

Hart's Tongue.

In shady lanes under hollow ways, and on shady walls. Found against the North side of the steeple of Loughton Church; and in the brick work of a cellar window at Mr. Foulis's, Woodford Row: not very common.

PILOSELLA repens. *Raii Syn.* 170.

HIERACIUM [*Pilosella*] foliis integerrimis ovatis
subtus tomentosis, caule repente, scapo uni-
floro. *Hudsoni Fl.* 298.

Classis Linnæi Syngenesia: Polygamia Æqualis.

Common creeping Mouse-ear.

In fields on a dry soil common.

It flowers in May.

PIMPINELLA saxifraga minor. *Raii Syn.* 213.

PIMPINELLA [*saxifraga*] foliis pinnatis, foliolis
radicalibus subrotundis: summis linearibus.

Hudsoni Fl. 111. β.

Classis Linnæi Pentandria Digynia.

Small Burnet Saxifrage, with divided leaves.

*In fields, on a dry gravelly soil: not very
common.*

It flowers in August.

PIMPINELLA saxifraga minor foliis sangui-
forbæ. *Raii Syn.* 213.

PIMPINELLA [*saxifraga*] foliis pinnatis, foliolis
radicalibus subrotundis: summis linearibus.

Hudsoni Fl. 111.

Classis Linnæi Pentandria Digynia.

The lesser round leaved Burnet Saxifrage.

*In fields, on a dry, gravelly soil: not very
uncommon.*

It flowers in August.

PINUS sylvestris foliis brevibus glaucis, conis parvis albertibus. *Raii Syn.* 442.

PINUS [*sylvestris*] foliis geminis: primordialibus solitariis glabris. *Hudsoni Fl.* 361.

Classis Linnæi Monoecia Monadelphica.

The Scotch Fir.

Found self-sown, near the great Pond at Snaresbrook: uncommon.

It flowers in May.

PLANTAGO aquatica. *Raii Syn.* 257.

ALISMA [*Plantago.*] foliis ovatis acutis, fructibus obtusis trigonis. *Hudsoni Fl.* 137.

Classis Linnæi Hexandria Polygynia.

Great Water Plantain.

In moist places, and in the sides of ponds and rivers: very common.

It flowers in July or August.

PLANTAGO foliis laciniatis, Coronopus dicta, Coronopus vulgaris, seu Cornu cervinum. *Raii Syn.* 315.

PLANTAGO [*Coronopus*] foliis linearibus pinnato-dentatis, scapo tereti. *Hudsoni Fl.* 52.

Classis Linnæi Tetrandia Monogynia.

Buckthorn Plantain. Star of the Earth.

On dry gravelly land. Found in open parts of the Forest, by the road side: not very common.

It flowers in July or August.

PLANTAGO latifolia vulgaris. *Raii Syn.* 314.

PLANTAGO [*major*] foliis ovatis glabris, scapo tereti, spica flosculis imbricatis. *Hudsoni Fl.* 51.

Classis Linnæi Tetrandia Monogynia.

Great Plantain, or Waybread.

In fields, and by the road side : very common.

It flowers in June or July.

PLANTAGO major incana. *Raii Syn.* 314.

PLANTAGO [*media*] foliis ovato lanceolatis pubescentibus, spica cylindrica, scapo tereti. *Hudsoni Fl.* 51.

Classis Linnæi Tetrandia Monogynia.

Hoary Plantain.

On dry gravelly soil, and by the road side : very common.

It flowers in July or August.

PLANTAGO quinquenervia. *Raii Syn.* 314.

PLANTAGO [*lanceolata*] foliis lanceolatis, spica subovata nuda, scapo angulato. *Hudsoni Fl.* 52.

Classis Linnæi Tetrandia Monogynia.

Ribwort, or Ribwort Plantain.

In fields and meadows : very common.

It flowers in June, July or August.

POLYGALA. *Raii Syn.* * 287.

POLYGALA [*vulgaris*] floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis lineari-lanceolatis. *Hudsoni Fl.* 271.

Classis Linnæi Diadelphia Octandria.

Milk-wort.

In fields, on a dry soil; and on the Forest: common.

It flowers in May or June.

POLYGONUM brevi angustoque folio. *Raii Syn.* 146.

POLYGONUM [*aviculare*] floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. *Hudsoni Fl.* 149. β.

Classis Linnæi Octandria Trigynia.

Knot Grass, with a small and narrow leaf.

On gravelly soil, and by the road side: not very common.

It flowers from June to September.

POLYGONUM mas vulgare. *Raii Syn.* 146.

POLYGONUM [*aviculare*] floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. *Hudsoni Fl.* 149.

Classis Linnæi Octandria Trigynia.

Common Knot-grass.

In ground uncultivated; by the road and waysides: very common.

It flowers from June to October.

WOODFORDIENSES. 127

POLYGONUM oblongo angustoque folio. *Raii Syn.* 146.

POLYGONUM [*aviculare* γ] floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. *Hudsoni Fl.* 149.

Classis Linnæi Octandria Trigynia.

Knot Grass, with a more oblong, and narrow leaf.

On gravelly soil, and by the road side ; also among corn.

It flowers from June to September.

POLYPODIUM. *Raii Syn.* 117.

POLYPODIUM [*vulgare*] frondibus pinnatifidis : pinnis oblongis subserratis, obtusis, radice squamata. *Hudsoni Fl.* 387.

Classis Linnæi Cryptogamia : Filices.

Common Polypody.

On the banks of hollow ways ; on walls, and at the roots of trees, sometimes in holes of hollow oaks. Found on an oak tree on the Forest between Muncombe and the Bald-faced Stag.

POPULAGO. Sive Caltha Palustris. *Raii Syn.* 272.

CALTHA [*palustris*] *Hudsoni Fl.* 214.

Classis Linnæi Polyandria Polygynia.

Marsh-Marigold.

In moist places, and on the sides of rivers. Found in the fields near the river Rhodon, opposite the Eight Mile Stone in the Woodford-bridge road ; and in Hackney-marsh, near Lee Bridge : not uncommon.

It flowers in April.

POPULUS alba. *Raii Syn.* 446.

POPULUS [*alba*] foliis subrotundis dentato-angulatis, subtus tomentosis. *Hudsoni Fl.* 370.

Classis Linnæi Dioecia Octandria.

The White Poplar or Abele Tree.

In moist boggy places, as the black poplar.

It flowers in March.

POPULUS Libyca Plinii, *Kenp's Theophrasti.*

Raii Syn. 446.

POPULUS [*tremula*] foliis subrotundis, dentato-angulatis, utrinque glabris. *Hudsoni Fl.* 370.

Classis Linnæi Dioecia Octandria.

The Asp, Aspen or Trembling Poplar.

In moist woods. Found on the Forest near Earl Tylney's Park.

It flowers in March.

POPULUS nigra. *Raii Syn.* 446.

POPULUS [*nigra*] foliis deltoidibus acuminatis ferratis *Hudsoni Fl.* 370.

Classis Linnæi Dioecia Octandria.

The Black Poplar.

In moist boggy places, and by the sides of rivers. Found in Mob's Hole, between the Eagle at Snaresbrook, and the Woodford-bridge Road: and near the river Rhodon: not uncommon.

It flowers in March.

PORTULA. *Raii Syn.* 368.

PEPLIS [*Portulaca.*] *Hudsoni Fl.* 128.

Classis Linnæi Hexandria Monogynia.

Water Purlane.

In moist places, and where waters have been stagnated in winter : not uncommon, in most of the ponds on the Forest.

It flowers in September.

POTAMOGETON aquis immersum folio pellucido, lato, oblongo, acuto. *Raii Syn.* 148.

POTAMOGETON [*lucens*] foliis lanceolatis planis in petiolis desinentibus. *Hudsoni Fl.* 61.

Classis Linnæi Tetrandia Tetragynia.

Long leaved great Pond-weed, with pellucid leaves.

In ponds and still places in rivers : common.

It flowers in June.

POTAMOGETON foliis pennatis. *Raii Syn.* 150.

MYRIOPHYLLUM [*spicatum*] floribus masculis interruptè spicatis. *Hudsoni Fl.* 357.

Classis Linnæi Monoecia Polyandria.

Feathered or Spiked Water Mill-foil.

In ditches, and still waters : common.

It flowers in June or July.

POTAMOGITON seu Fontalis crispa. Tribulus aquaticus. *Raii Syn.* 149.

POTAMOGETON [*crispum*] foliis lanceolatis alternis undulatis ferratis. *Hudsoni Fl.* 61.

Classis Linnæi Tetrandria Tetragynia.

The greater Water Caltrops.

In ditches, stagnated waters, and rivers. Found in the river Rhodon : common.

It flowers in May or June.

POTOMAGITON seu Fontalis media lucens. *Raii Syn.* 149.

POTAMOGETON [*serratum*] foliis lanceolatis oppositis subundulatis. *Hudsoni Fl.* 61.

Classis Linnæi Tetrandria Tetragynia.

The lesser Water-caltrops, or Frog's Lettuce.

In rivers, where the current is not rapid : not uncommon.

It flowers in June.

POTOMAGITON millefolium, seu foliis gramineis ramosum. *Raii Syn.* 150.

POTOMAGETON [*pectinatum*] foliis cetaceis parallelis approximatis distichis. *Hudsoni Fl.* 62.

Classis Linnæi Tetrandria Tetragynia.

Fine, or Fennel-leaved Pond-weed.

In large ponds and rivers. Found in the river Rhodon : not uncommon.

It flowers in June.

WOODFORDIENSES. 131

POTOMAGITON perfoliatum. *Raii Syn.* 149.

POTAMOGETON [*perfoliatum*] foliis cordatis amplexicaulibus. *Hudsoni Fl.* 60.

Classis Linnæi Tetrandria Trigynia.

Perfoliated Pond-weed.

In ponds and rivers. Found in the river Rhodon: common.

It flowers in June or July.

POTAMOGETON rotundifolium. *Raii Syn.* 148.

POTAMOGETON [*natans*] foliis oblongo-ovatis petiolatis natantibus. *Hudsoni Fl.* 60.

Classis Linnæi Tetrandria Trigynia.

Broad leaved Pond-weed.

In fish-ponds, and standing waters. Found in small rivers: very common.

It flowers in August.

PRIMULA pratensis inodora lutea. *Raii Syn.* 284.

PRIMULA [*veris*] foliis dentatis rugosis, scapho multifloro. *Hudsoni Fl.* 70. β.

Classis Linnæi Pentandria Monogynia.

Great Cowslips or Oxlips.

In woods, under hedges, and in meadows. Found in a meadow near Wray-House, Sir James Wright's, adjoining to the river Rhodon: uncommon.

It flowers in April or May.

PRIMULA veris major. *Raii Syn.* 284.

PRIMULA [*veris*] foliis dentatis rugosis, scapo multifloro. *Hudsoni Fl.* 70.

Classis Linnæi Pentandria Monogynia.

Common Pægils, or Cowslips.

In meadows : very common.

It flowers in April or May.

PRIMULA veris vulgaris. *Raii Syn.* 284.

PRIMULA [*vulgaris*] foliis dentatis rugosis, scapho unifloro. *Hudsoni Fl.* 70.

Classis Linnæi Pentandria Monogynia.

Common Primrose.

In woods, on the sides of fields, and under hedges : very common.

It flowers in April or May.

PRUNELLA. *Raii Syn.* 238.

PRUNELLA [*vulgaris*] foliis ovato-oblongis petiolatis. *Hudsoni Fl.* 231.

Classis Linnæi Didynamia Gymnospermia.

Common Self-heal.

In fields and meadows : very common. Found with a white flower ; but not common.

It flowers in August.

PRUNUS sylvestris. *Raii Syn.* 461.

PRUNUS [*spinosa*] spinosa, foliis lanceolatis pedunculis solitariis. *Hudsoni Fl.* 186.

Classis Linnæi Icosandria Monogynia.

The Black Thorn, or Sloe Tree.

In hedges: very common.

It flowers in March and April.

PRUNUS sylvestris major. *Raii Syn.* 462.

PRUNUS [*insititia*] spinosa, foliis ovatis subtus villosis, pedunculis geminis. *Hudsoni Fl.* 186. β.

Classis Linnæi Icosandria Monogynia.

The Black Bullace Tree.

In hedges. Found in a lane between the new road from Lee Bridge, in Marsh-street, Walthamstow; and in the woody part of the Forest, between The Windmill and Laytonstone: not very common.

It flowers in April.

PRUNUS sylvestris fructu majore albo. *Raii Syn.* 462.

PRUNUS [*insititia*] spinosa, foliis ovatis subtus villosis, pedunculis geminis. *Hudsoni Fl.* 186.

Classis Linnæi Icosandria Monogynia.

The White Bullace Tree.

In hedges. Found in the woody part of the Forest, between The Windmill and Laytonstone: not common.

It flowers in April.

PTARMICA. *Raii Syn.* 183.

ACHILLEA [*Ptarmica*] foliis lanceolatis, acuminatis argute ferratis. *Hudsoni Fl.* 325.

Classis Linnæi Syngenefia: Polygamia Superflua.

Sneeze wort. Bastard Pellitory, or Goose Tongue.

In meadows, and moist places on the Forest: not uncommon.

It flowers in August.

PULEGIUM. *Raii Syn.* 235.

MENTHA [*Pulegium*] floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus. *Hudsoni Fl.* 224.

Classis Linnæi Didynamia Gymnospermia.

Penny-royal, or Pudding-grafs.

In marshy moist places, where waters have stagnated in winter. Found in plenty on the hill before Mr. Warner's House, at Woodford Row, and about his Gravel Pit Pond: common.

It flowers in August or September.

PYRASTER seu *Pyrus sylvestris.* *Raii Syn.* 452.

PYRUS [*communis*] foliis ferratis, floribus corymbosis. *Hudsoni Fl.* 189.

Classis Linnæi Icosandria Pentagynia.

The wild Pear Tree.

In woods and hedges. Found in a hedge by the side of the river Rhodon, not far from Sir Edward Walpole's, Luxborough House.

It flowers in April.

Q.

QUERCUS latifolia. *Raii Syn.* 440.

QUERCUS [*Robur*] foliis deciduis oblongis superne latioribus: sinubus acutioribus; angulis obtusis. *Hudsoni Fl.* 359.

Classis Linnæi Monoecia Polyandria.

The common Oak Tree.

In woods, the woody parts of the Forest, and in hedges: common.

It flowers in April.

QUERCUS latifolia mas, quæ brevi pediculo est. *Raii Syn.* 440.

QUERCUS [*Robur*] foliis deciduis oblongis, superne latioribus, sinubus acutioribus, angulis obtusis. *Hudsoni Fl.* 359.

Classis Linnæi Monoecia Polyandria.

The Oak Tree with the Acorns on short footstalks.

As the former. Found on the Forest between Muncombe and The Bald-faced Stag: not common.

It flowers in April.

R.

RADICULA sylvestris feu palustris. *Raii Syn.*

301.

SISYMBRIUM [*amphibium*] filiquis declinatis oblongo-ovatis, foliis pinnatifidis ferratis.

Hudsoni Fl. 258. α.

Classis Linnæi Tetradynamia Siliquosa.

Water Radish.

In moist places, running waters, and by the sides of rivers. In the river Rhodon, near Luxborough House, Sir Edward Walpole's: common.

It flowers in June.

RADIOLA vulgaris serpyllifolia. Mille grana minima. *Raii Syn.* 345.

LINUM [*Radiola*] foliis oppositis, caule dichotomo, floribus tetrandis tetragynis. *Hudsoni Fl.* 117.

Classis Linnæi Pentandria Pentagynia.

The least Rupture-wort or All-feed.

In moist ground on sandy heaths. Found in a boggy soil, a little way in the wood, opposite The Bald-faced Stag: not common.

It flowers in August.

RANUNCULO five Polyanthemo aquatili albo affine. *Raii Syn.* 250.

RANUNCULUS [*aquatilis* γ] foliis omnibus capillaceis, circumscriptione oblongis. *Hudsoni Fl.* 213.

Classis Linnæi Polyandria Polygynia.

Fennel leaved Water Crowfoot.

In running waters. Found in the river Rhodon in plenty.

It flowers from April to June.

RANUNCULUS aquaticus albus, circinatis tenuissime divisis foliis, floribus ex alis longis pediculis innixis. *Raii Syn.* 249.

RANUNCULUS [*aquatilis* δ] foliis omnibus capillaceis circumscriptione rotundis. *Hudsoni Fl.* 213.

Classis Linnæi Polyandria Polygynia.

Finest leaved Water Crowfoot.

In deep stagnated waters: common.

It flowers from April to June.

RANUNCULUS aquatilis. *Raii Syn.* 249.

RANUNCULUS [*aquatilis*] foliis submersis capillaceis, emersis peltatis. *Hudsoni Fl.* 213.

Classis Linnæi Polyandria Polygynia.

Various water-leaved Crowfoot.

In ditches and standing waters; and in rivers where the current is not strong: common.

It flowers in April and May.

RANUNCULUS aquatilis hederaceus albus. *Raii Syn.* 249.

RANUNCULUS [*hederaceus*] foliis subrotundis trilobis integerrimis, caule repente. *Hudsoni Fl.* 212.

Classis Linnæi Polyandria Polygynia.

Ivy leaved Water Crowfoot.

In ditches, and in still or shallow parts of rivers, or ponds about the Forest: not common.

It flowers in May.

RANUNCULUS bulbosus. *Raii Syn.* 247.

RANUNCULUS [*bulbosus*] calycibus retroflexis, pedunculis fulcatis, caule erecto, foliis compositis. *Hudsoni Fl.* 211.

Classis Linnæi Polyandria Polygynia.

Round rooted or bulbous Crowfoot. Butter cups.

In fields and meadows: very common.

It flowers in May.

RANUNCULUS aquatilis omnino tenuifolius. *Raii Syn.* 249.

RANUNCULUS [*β aquatilis*] foliis capillaceis circumscriptione vaga brevioribus. *Hudsoni Fl.* 213. *β.*

Classis Linnæi Polyandria Polygynia.

Fine leaved Water Crowfoot.

In shallow ponds, usually where the water has been dried up: common.

It flowers from April to June.

RANUNCULUS arvorum. *Raii Syn.* 248.

RANUNCULUS [*arvensis*] feminibus aculeatis,
foliis superioribus decompositis linearibus.

Hudsoni Fl. 212.

Classis Linnæi Polyandria Polygynia.

Corn Crowfoot.

In fields, chiefly among corn.

It flowers in June.

RANUNCULUS flammeus minor. *Raii Syn.* 250.

RANUNCULUS [*flammula*] foliis ovato lanceo-
latis petiolatis, caule declinato. *Hudsoni Fl.*

210.

Classis Linnæi Polyandria Polygynia.

The lesser Spearwort.

*In moist meadows, and where waters have
stagnated in winter : very common.*

It flowers from June to September.

RANUNCULUS nemorosus dulcis, secundus Tragi.

Raii Syn. 248.

RANUNCULUS [*auricomus*] foliis radicalibus
reniformibus crenatis incis: caulinis digi-
tatis linearibus, caule multifloro. *Hudsoni*

Fl. 211.

Classis Linnæi Polyandria Polygynia.

Sweet Wood Crowfoot or Goldilocks.

In woods and under hedges : common.

It flowers in April.

RANUNCULUS palustris. *Raii Syn.* 249.

RANUNCULUS [*sceleratus*] foliis inferioribus palmatis, summis digitatis, fructibus oblongis.

Hudsoni Fl. 212.

Classis Linnæi Polyandria Polygynia.

Round leaved Water Crowfoot.

By the sides of ponds, and in moist ditches : common.

It flowers in May or June.

RANUNCULUS pratensis erectus acris. *Raii Syn.* 248.

RANUNCULUS [*acris*] calycibus patulis, pedunculis teretibus, foliis tripartito multifidis; summis linearibus. *Hudsoni Fl.* 211.

Classis Linnæi Polyandria Polygynia.

Upright Meadow Crowfoot.

In fields and meadows : very common.

It flowers in June or July.

RANUNCULUS pratensis repens. *Raii Syn.* 247.

RANUNCULUS [*repens*] calycibus patulis, pedunculis fulcatis, stolonibus repentibus, foliis compositis. *Hudsoni Fl.* 210.

Classis Linnæi Polyandria Polygynia.

Common creeping Crowfoot, or Butter-cups.

In moist fields and meadows. Found chiefly on banks or ditches, under hedges : not uncommon.

It flowers in May.

RAPHANISTRUM filiqua articulata glabra majore et minore. *Raii Syn.* 296.

[RHAPHANUS] [*Raphanistrum*] filiquis teretibus, articulatis, lævibus, unilocularibus. *Hudsoni Fl.* 252.

Classis Linnæi Tetradynamia Siliquosa.

White flowered Charlock, with a jointed Cod.

In fields and among corn: very common.

It flowers in June or July.

RAPHANUS rusticanus. *Raii Syn.* 301.

COCHLEARIA [*armoracia*] foliis radicalibus lanceolatis, crenatis, caulinis incis. *Hudsoni Fl.* 248.

Classis Linnæi Tetradynamia Siliculosa.

Horse-radish.

In ditches, and by the sides of waters. Found on the banks of the river Rhodon, near Luxborough House, Sir Edward Walpole's.

It flowers in May.

RAPHANUS aquaticus foliis in profundas lacinias divis. *Raii Syn.* 301.

SISYMBRIUM [*amphibium*] filiquis declinatis oblongo-ovatis, foliis pinnatifidis ferratis. *Hudsoni Fl.* 258.

Classis Linnæi Tetradynamia Siliquosa.

Water Radish.

In running waters, and in rivers; in moist watery places, and by the sides of ditches: common.

It flowers in June.

RAPISTRUM arvorum. *Raii Syn.* 295.

SINAPIS [*arvensis*] filiquis multangulis torosoturgidis, rostro longioribus. *Hudsoni Fl.* 260.

Classis Linnæi Tetradynamia Siliquosa.

Charlock or Wild Mustard.

In fields, and among corn: very common.

It flowers in May.

RAPISTRUM flore luteo, filiqua glabra articulata. *Raii Syn.* 296.

RAPHANUS [*Raphanistrum* β] filiquis teretibus, articulatis, lævibus, unilocularibus. *Hudsoni Fl.* 252.

Classis Linnæi Tetradynamia Siliquosa.

Yellow flowered Charlock with a jointed cod.

In fields and among corn: very common.

It flowers in June or July.

RESEDA vulgaris. *Raii Syn.* 366.

RESEDA [*lutea*] foliis omnibus trifidis, inferioribus pinnatis. *Hudsoni Fl.* 181.

Classis Linnæi Dodecandria Trigynia.

Base Rocket.

In gravelly fields, and sometimes on walls. Found on a flint wall, in Mr. Warner's garden: not common.

It flowers in July.

RHAMNUS catharticus. *Raii Syn.* 466.

RHAMNUS [*catharticus*] spinis terminalibus, floribus quadrifidis dioicis. *Hudsoni Fl.* 83.

Classis Linnæi Pentandria Monogynia.

Buckthorn : or common Purging Thorn.

In woods and hedges. Found in a hedge by the side of the river Rhodon, not far from Luxborough House, Sir Edward Walpole's : and in the Woodford Bridge Road, near The Old Mill.

It flowers in April or May.

ROS folis folio rotundo. *Raii Syn.* 356.

DROSERA [*rotundifolia*] scapis radicatis, foliis orbiculatis. *Hudsoni Fl.* 117.

Classis Linnæi Pentandria Pentagynia.

Rosa-folis, or Round leaved Sun-dew.

In moist places and bogs. Found on a boggy part of the Forest, a little way in the wood, opposite The bald-faced Stag : not common.

It flowers in July or August.

ROSA fylvestris altera minor, flore albo nostras. *Raii Syn.* 455.

ROSA [*arvensis*] germinibus globosis, pedunculisque glabris, caule petiolisque aculeatis, floribus subcymosis. *Hudsoni Fl.* 192.

Classis Linnæi Icosandria Polygynia.

White flowered Dog's rose.

In hedges : very common.

It flowers in May or June

ROSA sylvestris inodora seu canina. *Raii Syn.* 454.

ROSA [*canina*] germinibus ovatis, pedunculisque glabris, caule petiolisque aculeatis. *Hudsoni Fl.* 192.

Classis Linnæi Icosandria Polygynia.

Common red flowered wild Dog's rose, or Hoptree.

In hedges: very common.

It flowers in May or June.

ROSA sylvestris odora. *Raii Syn.* 454.

ROSA [*Eglanteria*] germinibus globosis, pedunculisque glabris, caule aculeis sparsis rectis, petiolis scabris, foliolis acutis. *Hudsoni Fl.* 191.

Classis Linnæi Icosandria Polygynia.

The Sweet Briar or Eglantine.

In hedges. Found in a lane which leads from Chigwell to Chigwell Row: not common.

It flowers in May or June.

RUBUS major fructu nigro. *Raii Syn.* 467.

RUBUS [*fruticosus*] foliis quinato-digitatis ternatisque, caule petiolisque aculeatis. *Hudsoni Fl.* 193.

Classis Linnæi Icosandria Polygynia.

The common Bramble, or Black-berry Bush.

In woods and Hedges: very common.

It flowers from May to September.

RUBUS minor fructu cæruleo. *Raii Syn.* 467.

RUBUS [*cæsius*] foliis ternatis subnudis: lateralibus bilobis, caule aculeato tereti. *Hudsoni Fl.* 193.

Classis Linnæi Icosandria Polygynia.

The small Bramble, or Dew-berry Bush.

In fields, and among corn; sometimes in woods, or in hedges: common.

It flowers in June or July.

RUSCUS. *Raii Syn.* 262.

RUSCUS [*aculeatus*] foliis suprà floriferis nudis. *Hudsoni Fl.* 373.

Classis Linnæi Dioecia Syngenesia.

Knee Holly, or Butcher's Broom.

In the woody part of the Forest. Found near Woodford Row, in great plenty: common.

It flowers in March or April.

UTA MURARIA. *Raii Syn.* 122.

ASPENIUM [*Ruta muraria*] frondibus alternatim decompositis: foliis cuneiformibus arenulatis. *Hudsoni Fl.* 386.

Classis Linnæi Cryptogamia: Filices.

White Maiden-hair. Wall-rue. Tentwort.

On walls, and on banks in hollow ways. Found on an old wall at Layton-stone, near The Green-Man; and on the wall before Mr. Bosanquet's House on the Forest, between that place and Woodford.

S.

SAGITTA. *Raii Syn.* 258.

SAGITTARIA [*Sagittifolia*] foliis sagittatis acutis.

Hudsoni Fl. 358.

Classis Linnæi Monoecia Polyandria.

Arrow Head.

In ditches, ponds, and rivers. Found on the side of the river Rhodon, near Luxborough House, Sir Edward Walpole's: not uncommon.

It flowers in June.

SALICARIA vulgaris purpurea foliis oblongis.

Raii Syn. 367.

LITHRUM [*Salicaria*] foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris.

Hudsoni Fl. 179.

Classis Linnæi Dodecandria Monogynia.

Purple spiked Loose-strife, or Willow-herb.

In marshy ground, and on the side of ditches and rivers. Found in a ditch the lower end of Marsh-street, Walthamstow, near Hackney-Marsh, and on the sides of most parts of the river Rhodon: common.

It flowers in August or September.

SALIX. *Raii Syn.* 447.

SALIX [*alba*] foliis lanceolatis acuminatis serratis utrinque pubescentibus: serraturis infimis glandulosis. *Hudsoni Fl.* 366.

Classis Linnæi Dioecia Diandria.

The most common White Willow.

In woods and hedges, chiefly in moist places: very common.

It flowers in April.

SALIX folio articulato splendente flexilis. *Raii Syn.* 448.

SALIX [*amygdalina*] foliis serratis glabris, lanceolatis petiolatis, stipulis trapeziformibus. *Hudsoni Fl.* 363.

Classis Linnæi Dioecia Diandria.

The round eared shining Willow, or Almond leaved Willow.

In coppices, and on the banks of the rivers: common.

It flowers in May.

SALIX folio longissimo. *Raii Syn.* 450.

SALIX [*viminalis*] foliis subintegerrimis lanceolato-linearibus longissimis, acutis, subtus sericeis, ramis virgatis. *Hudsoni Fl.* 366.

Classis Linnæi Dioecia Diandria.

The Osier.

In moist places, and near the sides of ponds and rivers: very common.

It flowers in April.

SALIX latifolia rotunda. *Raii Syn.* 449.

SALIX [caprea] foliis ovatis rugosis subtus tomentosis undatis superne denticulatis. *Hudsoni Fl.* 366.

Classis Linnæi Dioecia Diandria.

The common round leaved Sallow.

In woods and hedges : very common.

It flowers in April or May.

SALIX pumila angustifolia, prona parte cinerea.
Raii Syn. 447.

SALIX [fusca] foliis integerrimis ovatis subtus villosis nitidis. *Hudsoni Fl.* 366.

Classis Linnæi Dioecia Diandria.

Creeping Dwarf Willow.

On heaths, and hilly ground. Found near The Windmill, and all the way from thence to the great pond at Snarebrook : common.

It flowers in May.

SAMBUCUS. *Raii Syn.* 461.

SAMBUCUS [nigra] cymis quinquepartitis, caule arboreo. *Hudsoni Fl.* 112.

Classis Linnæi Pentandria Trigynia.

Common Elder.

In hedges : very common.

It flowers in April.

WOODFORDIENSES. 149

SAMBUCUS humilis seu Ebulus. *Raii Syn.* 461.

SAMBUCUS [*Ebulus*] cymis tripartitis, stipulis foliaceis. *Hudsoni Fl.* 113.

Classis Linnæi Pentandria Trigynia.

Dwarf Elder, Walwort or Danewort.

Under hedges, and by the road side. Found in the Lane between Sewardstone, commonly called Sufan, and Waltham-Abbey: not common.

It flowers in July.

SANICULA five Diapensia. *Raii Syn.* 221.

SANICULA [*Europæa*] foliis radicalibus simplicibus, flosculis omnibus sessilibus. *Hudsoni Fl.* 96.

Classis Linnæi Pentandria Digynia.

Sanicle.

In woods and under hedges. Found in the road between Walthamslow and Lee Bridge: not very common.

It flowers in May or June.

SAXIFRAGA aurea. *Raii Syn.* 158.

CHRYSOSPLENIUM [*oppositifolium*] foliis oppositis. *Hudsoni Fl.* 156.

Classis Linnæi Decandria Digynia.

Common golden Saxifrage.

In moist ground in woods, and near springs: very common.

It flowers in April.

SAXIFRAGA verna annua humilior. *Raii Syn.*

354.

SAXIFRAGA [*trydactylites*] foliis cuneiformibus trifidis alternis, caule erecto ramofo. *Hudsoni Fl.* 159.

Classis Linnæi Decandria Digynia.

Rue leaved Sengreen, or, Rue Whitlow Gras.

On rubbish or old Walls. Found on moft, in and round about Woodford : common.

It flowers in April.

SCABIOSA major communior folio laciniato. *Raii Syn.* 191.

SCABIOSA [*arvensis*] corollulis quadrifidis radiantibus, caule hispido, foliis pinnatifidis hispidis. *Hudsoni Fl.* 50.

Classis Linnæi Terrandria Monogynia.

Common Field Scabious. Double flowered Scabious.

In fields, upon the sides of banks, and among corn : not uncommon.

It flowers in August.

SCABIOSA minor vulgaris. *Raii Syn.* 191.

SCABIOSA [*columbaria*] corollulis quinquefidis radiantibus ovatis crenatis, caulinis pinnatis fetaceis. *Hudsoni Fl.* 50.

Classis Linnæi Tetrandria Monogynia.

The lesser Field Scabious.

In dry fields, and on hilly places : not common.

It flowers in June or July.

WOODFORDIENSES. 151

SCABIOSA radice succisa, flore globofo. *Raii Syn.* 191.

SCABIOSA [*succisa*] corollulis quadrifidis æqualibus, caule simplici, ramis approximatis, foliis lanceolato ovatis. *Hudsoni Fl.* 50.

Classis Linnæi Tetrandria Monogynia.

Devil's Bit.

In fields, and upon the Forest: common.

It flowers in August or September.

SCANDIX femine rostrato vulgaris. *Raii Syn.* 207.

SCANDIX [*Pecten Veneris*] seminibus lævibus, rostro longissimo. *Hudsoni Fl.* 107.

Classis Linnæi Pentandria Digynia.

Shepherd's Needle, or Venus-comb.

In fields and among corn: common.

It flowers in June or July.

SCORODONIA seu salvia agrestis. *Raii Syn.* 245.

TEUCRIUM [*Scorodonia*] foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto. *Hudsoni Fl.* 219.

Classis Linnæi Didynamia Gymnospermia.

Wood Sage.

In woods, on forests, and under hedges: common.

It flowers in July.

SCROPHULARIA *aquatica* major. *Raii Syn.*
* 283.

SCROPHULARIA [*aquatica*] foliis cordatis petiolatis decurrentibus, caule membranis angulato, racemis terminalibus. *Hudsoni Fl.*
239.

Classis Linnæi Didynamia Angiospermia.

Water Figwort, or Water Betony.

In moist ditches, and on the banks of rivers: common.

It flowers in July.

SCROPHULARIA major. *Raii Syn.* * 283.

SCROPHULARIA [*nodosa*] foliis cordatis basi transversis, caule obtusangulo. *Hudsoni Fl.*
239.

Classis Linnæi Didynamia Angiospermia.

Common knobby rooted Figwort.

In woods, and under hedges, in moist land: common.

It flowers in July.

SEDUM minus hæmatoides. *Raii Syn.* 269.

SEDUM [*reflexum*] foliis subulatis sparsis, basi membranacea soluta: inferioribus recurvatis, floribus cymosis. *Hudsoni Fl.* 170.

Classis Linnæi Decandria Pentagynia.

Yellow Stonecrop, or Prick Madam.

On Walls, and on the tiling of houses: not uncommon.

It flowers in July.

WOODFORDIENSES. 153

SEDUM parvum acre flore luteo. *Raii Syn.* 270.

SEDUM [*acre*] foliis subovatis adnato-fessilibus gibbis erectiusculis alternis, cyma trifida. *Hudsoni Fl.* 171.

Classis Linnæi Decandria Pentagynia.

Wall Pepper, or Stonecrop.

On walls, and on the tiling of houses: not uncommon.

It flowers in June or July.

SEMPERVIVUM majus. *Raii Syn.* 269.

SEMPERVIVUM [*tectorum*] foliis ciliatis, propaginibus patentibus. *Hudsoni Fl.* 185.

Classis Linnæi Dodecandria Trigynia.

Houseleek, or Sengreen.

On walls, and on the tiling of houses: common.

It flowers in July.

SENECIO hirsutus viscidus major odoratus. *Raii Syn.* 178.

SENECIO [*viscosus*] corollis revolutis, foliis pinatifidis viscidis, squamis calycinis laxis longitudine perianthii. *Hudsoni Fl.* 316.

Classis Linnæi Syngenesia: Polygamia Superflua.

Cotton Groundsel, or strong scented Groundsel.

In sandy ground: not uncommon.

It flowers in June.

SENECIO minor latiore folio, five montana. *Raii Syn.* 178.

SENECIO [*sylvaticus*] corollis revolutis foliis pinnatifidis denticulatis, caule corymbofo erecto. *Hudsoni Fl.* 316.

Classis Linnæi Syngenesia: Polygamia Superflua.
Mountain Groundsel.

In woods and under hedges: not uncommon.

It flowers in July.

SENECIO vulgaris. *Raii Syn.* 178.

SENECIO [*vulgaris*] corollis nudis, foliis pin-
nato-sinuatis amplexicaulibus, floribus sparsis.
Hudsoni Fl. 315.

Classis Linnæi Syngenesia: Polygamia Superflua.
Common Groundsel or Simson.

*In fields and under hedges, as well as in cul-
tivated ground: very common.*

It flowers in May.

SERPILLUM vulgare. *Raii Syn.* 230.

THYMUS [*serpyllum*] floribus capitatis caulibus
repentibus foliis planis obtusis basi ciliatis.
Hudsoni Fl. 229.

Classis Linnæi Didynamia Gymnospermia.
Common Mother of Thyme.

On hilly dry ground: very common.

It flowers in July or August.

SERRATULA. *Raii Syn.* 196.

SERRATULA [*tinctoria*] foliis pinnatifidis : lacinia terminali maxima. *Hudsoni Fl.* 304.

Classis Linnæi Syngenesia : Polygamia Æqualis.

Saw-wort.

In woods and fields : not very uncommon.

It flowers in July.

SESELI pratense nostras. *Raii Syn.* 216.

SESELI [*Carnifolia*] foliis bipinnatis sublinearibus ; petiolis basi membranaceis, seminibus ovalibus. *Hudsoni Fl.* 106.

Classis Linnæi Pentandria Digynia.

Meadow Saxifrage.

In fields and moist meadows. Found in the fields between Woodford-bridge and Luxborough House, Sir Edward Walpole's : not uncommon.

It flowers in August.

SIDERITIS Anglica strumosa radice. *Raii Syn.* 242.

STACHYS [*palustris*] verticillis sex floris, foliis lineari lanceolatis, semiamplexicaulibus. *Hudsoni Fl.* 227.

Classis Linnæi Didynamia Gymnospermia.

Clown's All-heal.

In moist places, and near rivers : common.

It flowers in August.

SIDERITIS arvensis rubra. *Raii Syn.* 242.

GALEOPSIS [*Ladanum*] internodiis caulinis æqualibus: verticillis omnibus remotis. *Hudsoni Fl.* 225.

Classis Linnæi Didynamia Gymnospermia.

Narrow leaved All-heal, or Iron Wort.

In corn fields: common.

It flowers in June, July, or August.

SIDERITIS humilis lato obtuso folio. *Raii Syn.* 242.

GLECHOMA [*arvensis*] foliis cordatis oblongis crenatis, caule hispido erecto. *Hudsoni Fl.* 224.

Classis Linnæi Didynamia Gymnospermia.

Upright Ground Ivy.

In shady places and under hedges; in corn fields, and in all the ploughed lands near the river Rhodon: not uncommon.

It flowers in August.

SINAPI album filiqua hirsuta femine albo vel ruffo. *Raii Syn.* 295.

SINAPIS [*alba*] filiquis hispidis: rostro obliquo longissimo. *Hudsoni Fl.* 260.

Classis Linnæi Tetradynamia Siliquosa.

White Mustard.

Found by the sides of cultivated fields: not very common.

It flowers in August.

SINAPI fativum fecundum. *Raii Syn.* 295.

SINAPIS [*nigra*] filiquis glabris tetragonis.
Hudsoni Fl. 260.

Classis Linnæi Tetradynamia Siliquosa.

Common Mustard.

On the sides of fields, and the banks of ditches : common.

It flowers in June.

SISYMBRIUM Cardamine, feu Nasturtium aquaticum. *Raii Syn.* 300.

SISYMBRIUM [*Nasturtium*] filiquis declinatis, foliis pinnatis: foliolis subcordatis. *Hudsoni Fl.* 257.

Classis Linnæi Tetradynamia Siliquosa.

Water Cresses.

In moist places, on the side of moist ditches : common.

It flowers in June and July.

SISYMBRIUM hirsutum. *Raii Syn.* 233.

MENTHA [*aquatica*] floribus capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus. *Hudsoni Fl.* 223. γ.

Classis Linnæi Didynamia Gymnospermia.

Water-mint.

In moist places, and on the sides of rivers : not uncommon.

It flowers in July and August.

Sium aromaticum Sison Officinarum. *Raii Syn.*

211.

Sison [*Amomum*] foliis pinnatis, umbellis erectis.

Hudsoni Fl. 104.

Classis Linnæi Pentandria Digynia.

Bastard Stone Parsley.

In woods, and under moist hedges. Found in the foot-way, between Mr. Norris's, opposite the Eight Mile Stone, and Woodford Church : common.

It flowers in August or September.

Sium latifolium foliis variis. *Raii Syn.* 211.

Sium [*latifolium*] foliis pinnatis, umbella terminali. *Hudsoni Fl.* 103.

Classis Linnæi Pentandria Digynia.

Great Water Parsnep.

In marshy ground, and on the side of rivers. Found by the side of the river Rhodon : common.

It flowers in July or August.

Sium pusillum foliis variis. *Raii Syn.* 212.

Sison [*inundatum*] repens umbellis bifidis.

Hudsoni Fl. 104.

Classis Linnæi Pentandria Digynia.

The least Water Parsnep.

In ponds, and stagnated waters. Found in Mr. Warner's Gravel Pit at Woodford Row : common.

It flowers in May or June.

WOODFORDIENSES. 159

Sium umbellatum repens. *Raii Syn.* 211.

Sium [*nodiflorum*] foliis pinnatis umbellis axillaribus subsessilibus. *Hudsoni Fl.* 103.

Classis Linnæi Pentandria Digynia.

Creeping Water Parsnep.

In ditches, and on the banks of rivers: common.

It flowers in July.

Solanum lignosum seu *Dulcamara*. *Raii Syn.* 265.

Solanum [*Dulcamara*] caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis. *Hudsoni Fl.* 78.

Classis Linnæi Pentandria Monogynia.

Woody Nightshade, or Bitter Sweet.

In moist places under hedges: common.

It flowers in June or July.

Solanum pomo spinoso oblongo, flore calathoides, *Stramonium* vulgo dictum. *Raii Syn.* 266.

Datura [*Stramonium*] pericarpis spinosis erectis ovatis. *Hudsoni Fl.* 78.

Classis Linnæi Pentandria Monogynia.

Thorn Apple.

On dunghills, and in rich soils: not uncommon.

It flowers in August or September.

SOLANUM vulgare. *Raii Syn.* 265.

SOLANUM [*nigrum*] caule inermi herbaceo, foliis ovatis dentato angulatis, umbellis nutantibus. *Hudsoni Fl.* 78.

Classis Linnæi Pentandria Monogynia.

Common Nightshade.

On rubbish and dunghills ; sometimes in cultivated ground : common.

It flowers in June or July.

SONCHUS asper laciniatus. *Raii Syn.* 163.

SONCHUS [*oleraceus* δ] pedunculis tomentosis, calycibus lævibus. *Hudsoni Fl.* 294. γ .

Classis Linnæi Syngenesia : Polygamia Æqualis.

Prickly Sow Thistle with jagged leaves.

On rubbish, in fields, and in cultivated ground : very common.

It flowers from June to September.

SONCHUS lævis minor paucioribus laciniis. *Raii Syn.* 163.

SONCHUS [*oleraceus*] pedunculis tomentosis calycibus lævibus. *Hudsoni Fl.* 294. β .

Classis Linnæi Syngenesia Polygamia Æqualis.

Smooth Sow-thistle, with leaves less jagged.

On rubbish, in fields, and in cultivated ground : common.

It flowers from June to September.

WOODFORDIENSES. 161

SONCHUS asper non laciniatus. *Raii Syn.* 163.

SONCHUS [*oleraceus*] pedunculis tomentosis, calycibus lævibus. *Hudsoni Fl.* 294. δ.

Classis Linnæi Syngenefia : Polygamia Æqualis.

Prickly Sowthistle, with less jagged leaves.

In cultivated ground; in fields and on rubbish : common.

It flowers from June to August.

SONCHUS lævis. *Raii Syn.* 162.

SONCHUS [*oleraceus*] pedunculis tomentosis, calycibus lævibus. *Hudsoni Fl.* 294.

Classis Linnæi Syngenefia : Polygamia Æqualis.

Common Sowthistle.

On rubbish, in fields, and in cultivated ground : very common.

It flowers from June to September.

SONCHUS repens, multis Hieracium majus. *Raii Syn.* 163.

SONCHUS [*arvensis*] pedunculis calycibusque hispidis subumbellatis, foliis basi cordatis. *Hudsoni Fl.* 295.

Classis Linnæi Syngenefia : Polygamia Æqualis.

In fields, and under hedges : common.

Tree Sowthistle.

It flowers in August.

SORBUS sylvestris foliis domesticæ similis. *Raii*
Syn. 452.

SORBUS [*aucuparia*] foliis pinnatis, utrinque
 glabris. *Hudsoni Fl.* 189.

Classis Linnæi Icosandria Trigynia.

Quicken Tree. Roan Tree. Mountain Ash.

*In woods and hedges. Found in Mr. Warner's
 field; and between Woodford and Snarebrook :
 not common.*

It flowers in May.

SPARGANIUM non ramosum. *Raii Syn.* 437.

SPARGANIUM [*erectum*] foliis triquetris erectis.
Hudsoni Fl. 346. β .

Classis Linnæi Monoecia Triandria.

Bur-reed not branched.

*In ditches, marshy ground, and rivers : not
 uncommon.*

It flowers in July.

SPARGANIUM ramosum. *Raii Syn.* 437.

SPARGANIUM [*erectum*] foliis triquetris erectis.
Hudsoni Fl. 346.

Classis Linnæi Monoecia Triandria.

Great branched Bur-reed.

*In ditches, marshy ground, and rivers : not
 uncommon.*

It flowers in July.

WOODFORDIENSES. 163

SPERGULA purpurea. *Raii Syn.* 351.

ARENARIA [*rubra*] foliis linearibus, stipulis membranaceis vaginantibus. *Hudsoni Fl.* 169.

Classis Linnæi Decandria Trigynia.

Purple Spurrey. Purple flowered Chickweed,
In sandy dry ground: not very common.

It flowers in June or July.

SPHONDYLUM. *Raii Syn.* 205.

HERACLEUM [*Sphondylium*] foliolis pinnatifidis,
Hudsoni Fl. 102.

Classis Linnæi Pentandria Digynia.

Cow Parsnep.

On the sides of fields, and under hedges: common.

It flowers in July.

STELLARIA. *Raii Syn.* 289.

CALLITRICHE [*verna*] foliis superioribus ovalibus, floribus androgynis. *Hudsoni Fl.* 2.

Classis Linnæi Monandria Digynia.

Water-wort. Vernal Star-wort. Star-headed
Water Chickweed.

In ditches, and stagnated waters: common.

It flowers from March to October.

STELLARIA aquatica foliis longis tenuissimis.

Raii Syn. 290.

CALLATRICHE [*autumnalis*] foliis omnibus linearibus, apice bifidis, floribus hermaphroditis. *Hudsoni Fl.* 2.

Classis Linnæi Monandria Digynia.

Autumnal Starwort.

In ditches and stagnated waters: not very common.

It flowers in September.

SYMPHYTUM magnum. *Raii Syn.* 230.

SYMPHYTUM [*officinale*] foliis ovato-lanceolatis decurrentibus. *Hudsoni Fl.* 68.

Classis Linnæi Pentandria Monogynia.

Comfrey.

In moist places, and on the sides of rivers: common.

It flowers in May.

T.

TAMNUS racemosa, flore minore luteo pallescente. *Raii Syn.* 262.

TAMUS [*communis*] foliis cordatis indivisis. *Hudsoni Fl.* 369.

Classis Linnæi Dioecia Hexandria.

Black Briony.

Under hedges, and among bushes : common.

It flowers in June.

TANACETUM. *Raii Syn.* 188.

TANACETUM [*vulgare*] foliis bipinnatis ferratis. *Hudsoni Fl.* 310.

Classis Linnæi Syngenesia: Polygamia Æqualis. Common Tanfy.

By the way side, and on the sides of fields : not uncommon.

It flowers in August.

TAXUS.

TAXUS. *Raii Syn.* 445.

TAXUS [*baccata*] foliis approximatis. *Hudsoni Fl.* 372.

Classis Linnæi Dioecia Monadelphia.

The Eugh or Yew Tree.

In forests, and sometimes in hedges. Found between the King's Oak and Epping, on the Forest, near the Fourteen Mile Stone, and in a hedge, between Mr. Warner's and Ribton Lane.

It flowers in March or April.

THALICTRUM seu Thalictrum majus. *Raii Syn.* 203.

THALICTRUM [*flavum*] caule folioso sulcato, panicula multiplici erectâ. *Hudsoni Fl.* 216.

Classis Linnæi Polyandria Polygynia.

Meadow Rue.

In moist meadows, and by the sides of rivers: not uncommon.

It flowers in June.

THLASPI vulgatus. *Raii Syn.* 305.

THLASPI [*campestre*] filiculis subrotundis, foliis sagittatis dentatis incanis. *Hudsoni Fl.* 245.

Classis Linnæi Tetradynamia Siliculosa.

Mithridate Mustard. Bastard Cresses.

In fields, on a dry sandy soil, and among corn.

It flowers in June or July.

TILIA folio minore. *Raii Syn.* 473.

TILIA [*Europæa*] floribus nectario destitutis.
Hudsoni Fl. 204. β.

Classis Linnæi Polyandria Monogynia.

The small leaved Lime, or Linden Tree; in
some Countries, called Baste.

*In woods and hedges. Found in a field, on the
right hand of the road, near Mr. Lloyd's, Friday
Hill House: not common, except in plantations.*

It flowers in July.

TILIA vulgaris platyphyllos. *Raii Syn.* 473.

TILIA [*Europæa*] floribus nectario destitutis.
Hudsoni Fl. 204.

Classis Linnæi Polyandria Monogynia.

The common Lime, Line or Linden Tree.

*In woods and hedges: not common, except in
plantations.*

It flowers in July.

TITHYMALUS characias Amygdaloides. *Raii
Syn.* 312.

EUPHORBIA [*Amygdaloides*] umbella multifida:
dichotoma, involucellis perfoliatis, orbicu-
latis, foliis obtusis. *Hudsoni Fl.* 184.

Classis Linnæi Dodecandria Trigynia.

Wood Spurge.

*Under hedges, and in woods. Found on the
Forest, near the top of Haggard Lane: not un-
common.*

It flowers in April.

TITHYMALUS verrucosus. *Raii Syn.* 312.

EUPHORBIA [*verrucosa*] umbella quinquefida :
dichotoma, involucellis cordatis, foliis lanceo-
latis ferrulatis pubescentibus, capsulis verru-
cosis. *Hudsoni Fl.* 183.

Classis Linnæi Dodecandria Trigynia.

Rough fruited Spurge.

*In fields and meadows. Found between the
lower end of Ribton Lane and Woodford Bridge :
not very common.*

It flowers in August.

TITHYMALUS helioscopius. *Raii Syn.* 313.

EUPHORBIA [*Helioscopia*] umbella quinquefida :
trifida : dichotoma, involucellis obovatis foliis
cuneiformibus ferratis. *Hudsoni Fl.* 183.

Classis Linnæi Dodecandria Trigynia.

Sun-spurge or Wart-wort.

*In Kitchen gardens, rich meadows, and on
dunghills : common.*

It flowers in July.

TITHYMALUS leptophyllos. *Raii Syn.* 313.

EUPHORBIA [*exigua*] umbella trifida : dichotoma, involucellis lanceolatis, foliis linearibus.
Hudsoni Fl. 182.

Classis Linnæi Dodecandria Trigynia.

Dwarf Spurge. Small annual Spurge.

*In Corn Fields : Found between the lower end
of Ribton Lane and Woodford Bridge : not
uncommon.*

It flowers in July.

WOODFORDIENSES. 169

TITHYMALUS parvus annuus, foliis subrotundis non crenatis, Peplus dictus. *Raii Syn.* 313.

EUPHORBIA [*Peplus*] umbella trifida; dichotoma, involucellis ovatis, foliis integerrimis obovatis petiolatis. *Hudsoni Fl.* 182.

Classis Linnæi Dodecandria Trigynia.

Petty Spurge.

In Kitchen gardens, and rich meadows: common.

It flowers in July.

TITHYMALUS platyphyllos Fuchsi. *Raii Syn.* 312.

EUPHORBIA [*platyphyllos*] umbella quinquefida: trifida: dichotoma: involucellis carina pilosis, foliis ferratis lanceolatis sessilibus. *Hudsoni Fl.* 184.

Classis Linnæi Dodecandria Trigynia.

Broad leaved Spurge.

In fields, and among corn. Found in the fields between Woodford Row and Woodford Bridge: not common.

It flowers in July.

TORMENTILLA. *Raii Syn.* 257.

TORMENTILLA [*erecta*] caule erecto, foliis sessilibus. *Hudsoni Fl.* 197.

Classis Linnæi Icosandria Polygynia.

Tormentil. Sept foil.

In dry fields, and on the Forest: very common.

It flowers in June.

TRAGOPOGON luteum. *Raii Syn.* 171.

TRAGOPOGON [*pratense*] calycibus corollæ radium æquantibus, foliis integris strictis. *Hudsoni Fl.* 293.

Classis Linnæi Syngenesia : Polygamia Æqualis.

Yellow Goat's Beard. Go to bed at Noon.

In fields and pastures. Found in Mr. Warner's fields : common.

It flowers in June.

TRICHOMANES. *Raii Syn.* 119.

ASPLENIUM [*Trichomanes*] frondibus pinnatis pinnis subrotundis crenatis. *Hudsoni Fl.* 385.

Classis Linnæi Cryptogamia : Filices.

English black Maiden-hair.

Found on some walls belonging to Mr. Warner : not common.

TRIFOLIUM album umbella filiquosa. *Raii Syn.*
Indiculus plantarum dubiarum.

TRIFOLIUM [*repens*, hybridum] capitulis umbellaribus, leguminibus, tetraspermis, caule ascendente. *Hudsoni Fl.* 283. β.

Classis Linnæi Diadelphia Decandria.

White Trefoil with a coddled Umbel.

On marshy soils. Found in the road from London to Woodford Bridge, between the Seven and Eight Mile Stone : not common.

It flowers from May to September.

WOODFORDIENSES. 171

TRIFOLIUM arvense humile spicatum, sive Lagopus. *Raii Syn.* 330.

TRIFOLIUM [*arvense*] spicis villōsis ovalibus, dentibus calycinis setaceis æqualibus. *Hudsoni Fl.* 285.

Classis Linnæi Diadelphia Decandria.

Hare's Foot, or Hare's Foot Trefoil.

In dry fields, and on sandy ground: not uncommon.

It flowers in July or August.

TRIFOLIUM cochleatum folio cordato maculato. *Raii Syn.* 333.

MEDICAGO [*Arabica*] pedunculis racemosis, leguminibus cochleatis spinosis, caule diffuso. *Hudsoni Fl.* 288.

Classis Linnæi Diadelphia Decandria.

Heart Trefoil, or Claver.

In fields, meadows, and among corn: not uncommon.

It flowers in May.

TRIFOLIUM fragiferum. *Raii Syn.* 329.

TRIFOLIUM [*fragiferum*] capitulis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. *Hudsoni Fl.* 286.

Classis Linnæi Diadelphia Decandria.

Strawberry Trefoil.

In fields and moist meadows: not uncommon.

It flowers in July or August.

TRIFOLIUM lupulinum alterum minus. *Raii Syn.* 330.

TRIFOLIUM [*procumbens*] spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. *Hudsoni Fl.* 287.

Classis Linnæi Diadelphia Decandria.

The lesser Hop Trefoil. Decumbent Trefoil.
In fields, meadows, and on the Forest: common.
It flowers from May to August.

TRIFOLIUM lupulinum minimum. *Raii Syn.* 331.

TRIFOLIUM [*filiforme*] spicis subimbricatis, vexillis deflexis persistentibus, calycibus pedicellatis, caulibus procumbentibus. *Hudsoni Fl.* 287.

Classis Linnæi Diadelphia Decandria.

Small Trefoil.

Found only on the dry parts of the Forest: not very common.

It flowers in May or June.

TRIFOLIUM luteum lupulinum. *Raii Syn.* 331.

MEDICAGO [*lupulina*] spicis ovalibus, leguminibus reniformibus monospermis, caule procumbente. *Hudsoni Fl.* 288.

Classis Linnæi Diadelphia Decandria.

Melilot Trefoil.

On a dry barren soil, in fields and on heaths: common.

It flowers from May to August.

TRIFOLIUM parvum hirsutum floribus parvis dilute purpureis in glomerulis mollioribus et oblongis, femine magno. *Raii Syn.* 329.

TRIFOLIUM [*striatum*] capitulis sessilibus sub-lateralibus ovatis, calycibus striatis rotundatis. *Hudsoni Fl.* 285.

Classis Linnæi Diadelphia Decandria.

Knotted Trefoil, with softer heads and large seed.

In dry fields, and on sandy ground: common.

It flowers in June.

TRIFOLIUM pratense album. *Raii Syn.* 327.

TRIFOLIUM [*repens*] capitulis umbellaribus, leguminibus tetraspermis, caule repente. *Hudsoni Fl.* 283.

Classis Linnæi Diadelphia Decandria.

White flowered Meadow Trefoil. Dutch Clover.

In fields and meadows: very common.

It flowers from May to September.

TRIFOLIUM pratense luteum capitulo lupuli, vel agrarium. *Raii Syn.* 330.

TRIFOLIUM [*agrarium*] spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, caule erecto. *Hudsoni Fl.* 286.

Classis Linnæi Diadelphia Decandria.

Hop Trefoil.

In sandy, or gravelly soils, and among corn: not uncommon.

It flowers in June.

TRIFOLIUM pratense purpureum. *Raii Syn.* 328.

TRIFOLIUM [*pratense*] spicis subvillosis cinctis stipulis oppositis membranaceis, corollis monopetalis. *Hudsoni Fl.* 284.

Classis Linnæi Diadelphia Decandria.

Common Purple, or Honey Suckle Trefoil.

In fields and meadows : common.

It flowers from May to September.

TRIFOLIUM purpureum majus, foliis longioribus et angustioribus, floribus saturatioribus. *Raii Syn.* 328.

TRIFOLIUM [*medium*] spicis villosis globosis terminalibus, corollis monopetalis foliolis lanceolatis subintegerrimis, caule erecto. *Hudsoni Fl.* 284.

Classis Linnæi Diadelphia Decandria.

Long leaved purple Trefoil with deeper coloured flowers.

In meadows, and under hedges, by the sides of fields : not uncommon.

It flowers in July.

TURRITIS. *Raii Syn.* 293.

TURRITIS [*glabra*] foliis radicalibus dentatis hispidis, caulinis integerrimis amplexicaulibus glabris. *Hudsoni Fl.* 254.

Classis Linnæi Tetradynamia Siliquosa.

Great Tower Mustard.

In sandy fields and on dry banks. Found on a dry bank in the lane which leads from High Beech to Waltham Abbey : not very common.

It flowers in May.

TUSSILAGO. *Raii Syn.* 173.

TUSSILAGO [*Farfara*] scapo imbricato unifloro, foliis subcordatis angulatis denticulatis. *Hudsoni Fl.* 315.

Classis Linnæi Syngenefia : Polygamia Superflua.
Common Coltsfoot.

In moist places, on the sides of ponds, and adjoining to watry places. Found in a piece of ground belonging to Mr. Forbes, near the Eight Mile Stone in Woodford : common.

It flowers in March or April.

TYPHA. *Raii Syn.* 436.

TYPHA [*latifolia*] foliis subensiformibus, spica mascula femineaque approximatis. *Hudsoni Fl.* 345.

Classis Linnæi Monoecia Triandria.

Great Cat's Tail, or Reed Mace.

On the banks of rivers, the sides of fish ponds, and in moist places : common.

It flowers in July.

V.

VALERIANA sylvestris major. *Raii Syn.* 200.

VALERIANA [*officinalis*] floribus triandris, foliis omnibus pinnatis. *Hudsoni Fl.* 12.

Classis Linnæi Triandria Monogynia.

Great wild Valerian.

In moist places, and near water; sometimes in woods and bushy places: not uncommon.

It flowers in June.

VALERIANA sylvestris minor. *Raii Syn.* 200.

VALERIANA [*dioica*] floribus triandris dioicis, foliis radicalibus ovatis, caulinis pinnatis, *Hudsoni Fl.* 12.

Classis Linnæi Triandria Monogynia.

Small wild or marsh Valerian.

In moist meadows. Found on the sides of Hackney Marsh., in the new road from Lee Bridge: not uncommon.

It flowers in June.

VALERIANA sylvestris seu palustris minor altera.
Raii Syn. 200.

VALERIANA [*dioica*] floribus triandris dioicis,
foliis radicalibus ovatis, caulinis pinnatis.
Hudsoni Fl. 12. β .

Classis Linnæi Triandria Monogynia.

The other small wild or marsh Valerian.

In moist meadows : not common.

It flowers in June.

VALERIANELLA arvensis præcox humilis, femine compresso, sive Lactuca agnina. *Raii Syn.* 201.

VALERIANA [*Locusta*] floribus triandris caule dichotomo, foliis linearibus. *Hudsoni Fl.* 12.

Classis Linnæi Triandria Monogynia.

Lamb's Lettuce, or Corn Sallet.

Among corn, in fields, on rich cultivated ground, and in gardens : common.

It flowers in April or May.

VALERIANELLÆ vulgaris species major, ferotina. *Raii Syn.* 201.

VALERIANA [*locusta*] floribus triandris, caule dichotomo, foliis linearibus. *Hudsoni Fl.* 13. β .

Classis Linnæi Triandria Monogynia.

Late flowering Lamb's Lettuce, or Corn Sallet.

In fields and among corn.

It flowers in June or July.

VERBASCUM mas latifolium luteum. *Raii Syn.* 287.

VERBASCUM [*Thapsus*] foliis decurrentibus utrinque tomentosis. *Hudsoni Fl.* 75.

Classis Linnæi Pentandria Monogynia.

Great White Mullein. High Taper. Cow's Lungwort.

In dry, sandy, barren places, on dry banks by the way side: common.

It flowers in July.

VERBASCUM nigrum flore parvo, apicibus purpureis. *Raii Syn.* 288.

VERBASCUM [*nigrum*] foliis cordato oblongis petiolatis. *Hudsoni Fl.* 76.

Classis Linnæi Pentandria Monogynia.

Sage leaved black Mullein.

On dry banks, by the road side. Found in a lane leading from Chigwell to Chigwell Row: not very common.

It flowers in July.

VERBENA vulgaris. *Raii Syn.* 236.

VERBENA [*officinalis*] tetrandra, spicis filiformibus paniculatis, foliis multifido-lanceolatis, caule solitario. *Hudsoni Fl.* 505.

Classis Linnæi Diandria Monogynia.

Vervain.

By the road side, near foot-paths, and houses: very common.

It flowers from July to September.

WOODFORDIENSES. 179

VERBESINA feu Cannabina aquatica flore minus pulchro, elatior et magis frequens. *Raii Syn.* 187.

BIDENS [*tripartita*] foliis trifidis, corollis flosculis, calycibus subfoliolis, feminibus erectis. *Hudsoni Fl.* 309.

Classis Linnæi Syngenesia : Polygamia Æqualis. Trifid Water Hemp Agrimony, with a divided leaf.

In marshy, watry places, and stagnated waters: common.

It flowers in July or August.

VERBESINA pulchriore flore luteo. *Raii Syn.* 187.

VERBESINA [*cernua*] foliis lanceolatis, amplexicaulibus, floribus cernuis, feminibus erectis. *Hudsoni Fl.* 309.

Classis Linnæi Syngenesia : Polygamia Æqualis. Water Hemp Agrimony, with an undivided leaf. Whole leaved Water Hemp Agrimony.

In ditches and marshy ground: common.

It flowers in July or August.

VERONICA aquatica angustifolia minor. *Raii Syn.* 280.

VERONICA [*scutellata*] racemis lateralibus, foliis ovatis, pedicellis pendulis, foliis linearibus integerrimis. *Hudsoni Fl.* 5.

Classis Linnæi Diandria Monogynia.

Narrow leaved Water Speedwell, or Brooklime.

In wet ditches and marshy ground: common.

It flowers in June.

VERONICA aquatica longifolia media. *Raii Syn.*
280.

VERONICA [*Anagallis*] racemis lateralibus, foliis lanceolatis ferratis, caule erecto. *Hudsoni Fl.* 5.

Classis Linnæi Diandria Monogynia.

The middle long-leaved Water Speedwell or Brook-lime.

In wet ditches, and marshy ground : common.

It flowers in July.

VERONICA aquatica rotundifolia, Becabunga dicta, minor. *Raii Syn.* 280.

VERONICA [*Becabunga*] racemis lateralibus, foliis ovatis planis, caule repente. *Hudsoni Fl.* 4.

Classis Linnæi Diandria Monogynia.

Common Brooklime.

In wet ditches, and small running waters : common.

It flowers in June.

VERONICA Chamædrys fylvestris dicta. *Raii Syn.* 281.

VERONICA [*Chamædrys*] floribus recemosis lateralibus, foliis ovatis rugosis dentatis sessilibus, caule debili. *Hudsoni Fl.* 5.

Classis Linnæi Diandria Monogynia.

Wild Germander.

In fields and meadows : very common.

It flowers in May.

WOODFORDIENSES. 181

VERONICA floribus singularibus, in oblongis pediculis, Chamædryfolia. *Raii Syn.* 279.

VERONICA [*agrestis*] floribus solitariis, foliis cordatis incisis petiolatis. *Hudsoni Fl.* 6.

Classis Linnæi Diandria Monogynia.

Germander-speedwell or Chickweed.

In fields, on rubbish, and in cultivated ground: common.

It flowers in May.

VERONICA flosculis singularibus, cauliculis adhærentibus. *Raii Syn.* 279.

VERONICA [*arvensis*] floribus solitariis, foliis cordatis incisis sessilibus. *Hudsoni Fl.* 6.

Classis Linnæi Diandria Monogynia.

Speedwell or Chickweed.

In fields, under hedges, and on old walls: common.

It flowers in May.

VERONICA flosculis singularibus Hederulæ folio, morsus gallinæ minor dicta. *Raii Syn.* 280.

VERONICA [*Hederifolia*] floribus solitariis, foliis cordatis planis quinquelobis. *Hudsoni Fl.* 6.

Classis Linnæi Diandria Monogynia.

Ivy leaved Speedwell, or small Henbit.

In fields, and sowed ground: not uncommon.

It flowers in April or May.

VERONICA mas supina et vulgatissima. *Raii Syn.* 281.

VERONICA [*officinalis*] spicis lateralibus pedunculatis, foliis oppositis, caule procumbente. *Hudsoni Fl.* 4.

Classis Linnæi Diandria Monogynia.

The Male Speedwell or Fluellin.

In barren, dry fields, and on heaths: common.

It flowers in May.

VERONICA pratensis minor, five Betonica Pauli. *Raii Syn.* 279.

VERONICA [*serpyllifolia*] racemo terminali subspicato, foliis ovatis, glabris crenatis. *Hudsoni Fl.* 4.

Classis Linnæi Diandria Monogynia.

Little smooth Speedwell, or Fluellin, or Paul's Betony.

In fields and meadows: not uncommon.

It flowers in May.

VICIA sepium perennis. *Raii Syn.* 320.

VICIA [*sepium*] leguminibus pedicellatis subquaternis erectis, foliis ovatis integerrimis: exterioribus decrefcentibus. *Hudsoni Fl.* 278.

Classis Linnæi Diadelphía Decandria.

Bush Vetch.

In bushy places, and in hedges: very common.

It flowers in May.

WOODFORDIENSES. 183

VICIA sylvestris five Cracca major. *Raii Syn.*
321.

VICIA [*Lathyroides*] leguminibus sessilibus foliariis erectis glabris foliolis senis: inferioribus obcordatis. *Hudsoni Fl.* 279.

Classis Linnæi Diadelphia Decandria.

Strangle Tare, or wild Vetch.

In dry fields, usually in sandy soils: common.

It flowers in May.

VINCA Pervinca minor. *Raii Syn.* 268.

VINCA [*minor*] caulibus procumbentibus, foliis lanceolato-ovatis. *Hudsoni Fl.* 77.

Classis Linnæi Pentandria Monogynia.

Periwinkle.

Under hedges and in woods. Found under the hedge on the left hand of the foot way from Woodford to Woodford Church, between Mr. Bosanquet's and Mr. Loney's; and on the Islands in Mr. Warner's gravel pit pond: not common.

It flowers in May.

VIOLA bicolor arvensis. *Raii Syn.* 366.

VIOLA [*tricolor*] caule triquetro diffuso, foliis oblongis incisis, stipulis dentato-pinnatis, lacinia suprema oblonga crenata petiolata. *Hudsoni Fl.* 331. β.

Classis Linnæi Syngenesia: Monogamia.

Pansies. Heart's-ease. Two Faces under a Hood.

In fields, and under hedges: common.

It flowers from May to September.

VIOLA *martia inodora Sylvestris. Raii Syn.*
364.

VIOLA [*canina*] caule demum adscendente foliis
oblongo cordatis. *Hudsoni Fl.* 331.

Classis Linnæi Syngenesia : Monogamia.

Wild or Dog's Violet.

Under hedges, and on the sides of fields : very common.

It flowers in March.

VIOLA *martia major hirsuta inodora. Raii Syn.*
365.

VIOLA [*hirta*] acaulis, foliis cordatis piloso-
hispidis. *Hudsoni Fl.* 330.

Classis Linnæi Syngenesia : Monogamia.

Violet with Throat-wort Leaves. Hairy
Violet.

In woods and under hedges : not very common.

It flowers in March.

VIOLA *martia purpurea. Raii Syn.* 364.

VIOLA [*odorata*] acaulis, foliis cordatis, stolo-
nibus reptantibus. *Hudsoni Fl.* 330.

Classis Linnæi Syngenesia : Monogamia.

Purple sweet Violet.

Under hedges and fields : very common.

It flowers in March or April.

VIOLA tricolor. *Raii Syn.* 361.

VIOLA [*tricolor*] caule triquetro diffuso, foliis oblongis incis, stipulis dentato-pinnatis, lacinia suprema oblonga crenata petiolata. *Hudsoni Fl.* 331.

Classis Linnæi Syngenefia Monogamia.

Panflies. Heart's-eafe. Three Faces under a Hood.

In fields and under hedges : common.

It flowers from May to September.

VIRGA AUREA. *Raii Syn.* 176.

SOLIDAGO [*Virga aurea*] caule subflexuoso angulato, racemis subpaniculatis erectis confertis. *Hudsoni Fl.* 318.

Classis Linnæi Syngenefia: Polygamia Superflua.

Common Golden Rod.

In woods, under hedges, and on heaths : not uncommon.

It flowers in August.

VISCUM. *Raii Syn.* 464.

VISCUM [*album*] foliis lanceolatis obtusis, caule dichotomo, spicis axillaribus. *Hudsoni Fl.*

367.

Classis Linnæi Monoecia Tetrandia.

Missel or Misseltoe.

On trees, particularly, the oak, apple, pear, ash, lime, willow, elm, &c. &c. Found on an oak, between Woodford Row and The Bald Faced Stag, near The Ten Mile Stone : and on an apple-tree in an Orchard in Loughton : and on several trees, many of them oaks, between that place and Mr. Conyers's, Copped Hall.

It flowers in May.

U.

ULMARIA. *Raii Syn.* 259.

SPIRÆA [*Ulmaria*] foliis pinnatis: impari majore lobato, caule herbaceo, floribus cymosis.
Hudsoni Fl. 191.

Classis Linnæi Icosandria Polygynia.

Meadow-sweet.

In moist meadows, and on the banks of rivers.

Found near the river Rhodon: common.

It flowers in June or July.

ULMUS folio glabro. *Raii Syn.* 469.

ULMUS [*compestris*] foliis ovatis duplicato-ferratis basi inæqualibus. *Hudsoni Fl.* 94. β.

Classis Linnæi Pentandria Digynia.

The smooth leaved or Wych Elm.

In hedges. Found before Mr. Norris's house, in Woodford, almost opposite the eight mile stone.

It flowers in April.

ULMUS folio latissimo scabro. *Raii Syn.* 469.

ULMUS [*glabra*] foliis oblongo-ovatis duplicato-ferratis basi inæqualibus, cortice glabro. *Hudsoni Fl.* 95.

Classis Linnæi Pentandria Digynia.

The Wych-Hasel, or broad leaved Elm.

In hedges and woods. Found in a copse belonging to Mr. Warner, adjoining to Ribton-lane: very uncommon.

It flowers in April.

ULMUS vulgatissima folio lato scabro. *Raii Syn.* 468.

ULMUS campestris foliis ovatis duplicato-ferratis basi inæqualibus. *Hudsoni Fl.* 94.

Classis Linnæi Pentandria Digynia.

Common Elm.

In hedges: very common.

It flowers in April.

URTICA minor. *Raii Syn.* 140.

URTICA [*urens*] foliis oppositis ovalibus. *Hudsoni Fl.* 355.

Classis Linnæi Monoecia Tetrandria.

The lesser Stinging Nettle.

In cultivated, and uncultivated ground: very common.

It flowers in August.

WOODFORDIENSES. 189.

URTICA racemifera major perennis. *Raii Syn.*
139.

URTICA [*dioica*] foliis oppositis cordatis, racemis geminis. *Hudsoni Fl.* 355.

Classis Linnæi Monoecia Tetrandria.

Common Stinging Nettle.

Under hedges, and on rubbish: very common.

It flowers in July.

A P P E N D I X.

BIFOLIUM majus seu *Ophris* major quibusdam.
Raii Syn. 385.

OPHRYS [*ovata*] bulbo fibroso, foliis ovatis,
nectarii labio bifido. *Hudsoni Fl.* 338.

Classis Linnæi Gynandria Diandria.

Common Tway blade.

In woods and among bushes, sometimes in meadows and fields. Found in a copse, the corner of Mr. Warner's large meadow; and on the Forest, between Woodford Row and Chingford Hatch: not common.

It flowers in June.

GALE frutex odoratus septentrionalium, *Elæagnus* Cordi. *Raii Syn.* 443.

MYRICA [*Gale*] foliis lanceolatis subserratis,
caule suffruticoso. *Hudsoni Fl.* 368.

Classis Linnæi Dioecia Tetrandria.

Goule. Sweet Willow. Dutch Myrtle.

In moist marshy places. Found near Mr. Warner's gravel-pit pond, at Woodford Row: very uncommon.

It flowers in May.

POLYGONATUM. *Raii Syn.* 263.

CONVALLARIA [*multiflora*] foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris. *Hudsoni Fl.* 126.

Classis Linnæi Hexandria Monogynia.

Solomon's Seal.

In woods. Found in a copse, the corner of Mr. Warner's large meadow: very uncommon.

It flowers in May or June.

I N D E X

O F T H E

E N G L I S H N A M E S.

A.

	Page
Abele Tree	128
Adder's Tongue	109
Agrimony	3
Dutch	50
Hemp	50
Alder, common	4
black, Berry bearing	55
All-good	17
Ale-hoof	22
Alkanet; bastard	19
All-heal, Clown's	155
narrow leaved	156
All-feed, upright Blite	34
least Rupture-wort	136
Angelica, wild	9
Apple-Thorn	159
Archangel, red	79
yellow	78
white	77
Arrow-head	146

C c

Arsmart,

	Page
Arsmart, dead or spotted	120
greater, with pale leaves	121
with hoary leaves	120
with spotted stalks	120
willow leaved, perennial, or	
water pepper	121
Ash Tree, common	55
mountain	162
Ash-weed	9
Asp, or Aspen	128
Avens, common	27

B.

Basil, great-wild	38
Bastard Cresses	166
Base Rocket	142
Baste	167
Batchelor's-buttons	91
Bear's-foot	64
Bed-straw, Ladies-yellow	57
Beech Tree	51
Bell-flower, lesser round leaved	23
Berberry	15
Bethwyne	37
Betony, water	152
wood	15
Bindweed black	51
great	38
small	38
Birch Tree	16
	Bird

Bird Cherry	31
Bird's Foot	112
Trefoil	88
greater	88
Bird's Nest	43
Bitter-sweet	159
Blackberry	144
Blite, late flowering, or Fig-leaved	16
the least	18
maple leaved	35
upright	34
round leaved, or all-feed	34
Borage	18
Bottle, blue	41
white	89
Box Tree	21
Brakes, common	52
Bramble, or Briar, common	144
small	145
Briar, sweet	144
Briony, black	165
white	18
Broom, common	58
rape	113
Brooklime, common	180
middle, long leaved	180
narrow leaved	179
Buckbean	97
Buckthorn	143
Buck's-horn plantain	124
Bugle	19
C c 2	Bugloss,

	Page
Bugloss, small wild	19
vipers	46
Bullace Tree, black	133
white	133
Burdock	82
with smaller heads	83
taller, with a purple flower	82
Burnet Saxifrage, lesser round leaved	123
small	123
Bur-reed, great branched	162
not branched	162
Bush Vetch	182
Butcher's Broom	145
Butter-bur, common	122
Butter cups	138
common creeping	140

C.

Calf's-snout	10
Caltrops, water, greater	130
lesser	130
Cammock	9
Camomile, sweet scented creeping	33
Campion-bladder	89
corn, white	89
red-flowered	91
wild, white	91
Canterbury bells	23
Carrot, wild	43
Cat-mint	105
Cat's-	

I N D E X.

197

Page

Car's-tail, great	175
Celandine, greater	115
leffer	34
Centaury, leffer	31
Charlock, white	141
or white Mustard	142
yellow	142
Cheefe-rening	57
Cheir, wild	85
Cherry, wild cluster, or bird	31
Cherry-wood	110
Chervil, small Hemlock	103
wild	32
Chesnut Tree	29
Chick-weed, breakstone	7
least	6
marsh, great	6
moufe-ear, broader-leaved	5
least	32
plantain leaved	7
purple flowered	163
water, long leaved	5
small	6
star-headed	163
Chickweed or Speedwell	181
Cicely, wild	36
Cinque-foil, common	118
Tormentil	118
Clary, common English wild	66
Claver	171
Climber, great wild	37
	Clot-

	Page
Clot-bur	82
finaller heads	83
taller, with a purple flower	82
woolly heads	83
Clover	173
Clown's All-heal	155
Cockle	90
Cock's-comb, great marsh, red rattled	117
red rattle	118
yellow	117
Cock's-head	109
Codlings and Cream	93
Colt's-foot, common	175
Comfrey	164
Corn, bottles	41
Feverfew	33
Marygold	35
Mint	98
Parsley, small	30
Rose	115
Sallet	177
late flowering	177
small	76
Violet, coddled	22
Cornel, female	39
Cotton Thistle	26
Cow's Lung-wort	178
Cow Parsnep	163
Cowslip, great	132
Cow-weed	36
Cow-wheat, common, wild	96
	Crab

I N D E X.

199

Page

Crab Tree	96
Crane's-bill, dove's-foot	60
field	59
hemlock leaved	59
jagged leaved	60
small flowered	60
Cresses, bastard	166
water	157
winter, or rocket	14
Crimson Grass Vetch	30
Crowfoot, common creeping	140
corn	139
finest leaved	137
ivy leaved	138
round, rooted, or bulbous	138
sweet-wood	139
Water fennel leaved	137
fine leaved	138
round leaved	140
various leaved	137
Crow-garlick	4
Cuckow flower	24
pint	11
Cudweed, black headed	61
common	62
least	61
long leaved, low branched	61

Daffodil,

D.

	Page
Daffodil, wild English	104
Daisie, common	15
greater	85
Dandelion	44
rough, commonly called Dandelion- Hawkweed	44
Deptford Pink	29
Devil's-bit	151
yellow	66
guts	41
Dewberry	145
Dock, common broad-leaved	82
greener leaved	81
sharp pointed	80
lesser	81
with curled leaves	81
great water	79
Dodder	41
Dogberry	39
Dog's Mercury	99
Rose, red flower	144
white flower	143
Violet	184
Dove's-foot	60
jagged-leaved	60
small flowered	60
Dropwort-hemlock	109
water	108
lesser	108
	Duck's

I N D E X.

201

Page

Dropwort, common	52
Duck's-meat	84
greater	85
ivy leaved	84
Dwale	14
Dwarf Elder	149
Dyer's-weed, or wild Woad	89
Wood-waxen	59

E.

Earth-nut	20
Eglantine	144
Elder, common	148
Dwarf	149
Elecampane	63
Elm, broad-leaved	188
common	188
smooth-leaved, or Wych	187
Enchanter's Nightshade	37
Eugh Tree	166
Eye-bright	50
red	50

F.

Fennel, common	54
Fern, common male	54
female, or common brakes	52
great branched, with indented leaves	53
lesser branched	53
male prickly auriculate	52
water, or flowering	53

D d

Fever-

	Page
Feverfew	96
corn	33
field	33
Figwort, common knobby rooted	152
water	152
Finckle	54
Five leaved grafs	118
Flag, sweet smelling or Calamus	2
Flax, purging or wild dwarf	87
Fleabane, middle	39
small	39
Flix-weed	49
Flower-de-luce, yellow water	73
Fluellin	182
sharp pointed	86
Fool's parsley	36
stones, female	111
male	110
Foxglove, purple	45
French Mercury, male and female	99
Frog's lettuce	130
Fumitory	56
greater ramping	56
Furze	59
leffer	58
needle	58

G.

	Page
Gatter-wood	39
Germander-Speedwell, or Chickweed	181
wild	180
Gill-go-by-the-ground	22
Gladdon or Gladwyn, stinking	73
Goat's-beard, yellow	170
Golden rod, common	185
Goldilocks	139
Goosefoot, common	17
sharp leaved	17
Goosegrafs	10
great	19
Goose Tongue	134
Gorse	59
lesser	58
Goule	190
Green-wood	59
Go-to-bed at noon	170
Gout-weed	9
Grass-vetch, crimson	30
Ground Ivy	22
upright	156
Groundfel, common	154
cotton, or strong scented	153
mountain	154

H.

	Page
Hard-beam Tree	114
Hare bells	67
Hare's foot	171
Hart's tongue	122
Hawk-nut, or rather Hog-nut	20
Hawkweed, bitten roots	66
bushy	65
long rooted	65
smooth succory	65
Hawthorn	100
Hazel-wych	188
-nut Tree	40
Heart's-ease	183
Three Faces under a Hood	185
Heath common, or Ling	48
cross leaved	47
fine leaved	48
Pea, or wood pea	113
Hedge Mustard	48
Nettle	57
Parsley	30
Hellebore, great bastard black	64
Hell-weed	41
Hemlock, Chervil small with rough seeds	103
dropwort	109
lesser, or Fool's parsley	36
water	122
Hemp-	

Hemp-agrimony, common or Dutch	50
Water, trifid	179
whole leaved	179
Henbane, common	68
Henbit, great	78
small	181
Hep-tree	144
Herb, Bennet, or common Avens	27
Gerard	9
Robert	61
Two-pence	105
High-taper	178
Hog-nut	20
Holly	3
Honeyfuckle, common	23
trefoil	174
Hop, male and female	89
trefoil	173
Horehound, stinking	14
water	91
Horn-beam Tree, sometimes Horse-beech	
or Horn-beech	114
Horse-radish	141
Horse-tail, corn	46
lesser, marsh	47
wood	47
Hound's-tongue, great	42
lesser green leaved	42
Houfeleek	153
Hyacinth, English	67

I.

	Page
Jack-by-the-hedge	64
Iron-wort	156
Ivy, climbing or berried	63
barren or creeping	63
Juniper, common	74

K.

Kipper-nut	20
Knapweed, common	71
great	71
Knawell, annual	75
Knee-holly	145
Knot-grass, common	126
German	75
oblong narrow leaved	127
small narrow leaved	126

L.

Ladies Bedstraw, mountain	101
white	101
yellow	57
Smock, common	24
Thistle	25
Traces, triple	112
Lamb's lettuce	177
late flowering	177
small	76
Lang-de-beuf	64
Larkspur,	

	Page
Marigold, corn	35
marsh	127
Marjoram, wild	112
Matfellow, common	71
gr et	71
May Lily	86
weed, stinking	32
Meadow Pink	90
Rue	166
Meadow-sweet	187
Medick-Vetchling	109
Melilot	97
trefoil	172
Mercury, common English	17
dog's	99
French, male and female	99
Milfoil	100
water, feathered or spiked	129
verticillated	119
Milkwort	126
Mill-mountain	87
Miltwaft	12
Mint, Nep or Nap	105
Pepper	98
Water	97
with whorled coronets	98
Missel or Misseltoe	186
Money-wort	105
purple flowered	106
Mother of thyme, common	154
Mountain Ash	162
	Moufe-

	Page
Mouse-ear	102
common creeping	123
tail	103
Mugwort	11
Mullein, great white	178
sage leaved black	178
Mustard, common	157
hedge	49
mithridate	166
tower, great	174
white	156

N.

Navew, wild	104
Needle Furze	58
Nep, or Nap or Cat-Mint	105
Nettle, dead	77
hemp-leaved, with a parti-co-	
loured flower	77
white flowered	77
red	79
yellow	78
stinging, common	189
lesser	188
Nettle-hemp	78
fair flowered	77
yellow	78
Nightshade, common	160
deadly	14
Enchanters	37
woody	159
E c	Nipple-

Nipple-wort	79
Nose-bleed	100
Nut Earth, Kipper, Pig, Hawk or Hog	20

O.

Oak Tree, common	135
with acorns on short foot-stalks	135
Oculus Christi, or Christ's eye	66
Orach, common wild	16
narrow leaved	12
with a spear pointed leaf	13
Orchis, butterfly lesser	110
female	111
female handed	111
male, or male fool's stones	110
male handed	111
Orpine	8
Osier	147
Osmund Royal	53
Ox-eye	85
lips	131
tongue	64

P.

Pagils	132
Pansies, two faces under a hood	183
three faces under a hood	185
Park-leaves	68
hairy	68
Parsley,	

I N D E X.

211

Page

Parsley, Corn small	30
hedge	30
knotted	31
Parsnep-Cow	163
great Water	158
least Water	158
water-creeping	159
wild	117
Paul's Betony	182
Pea Wood or Heath	113
Pear Tree, wild	134
Pellitory of the wall	116
Penny-royal	134
Penny-wort marsh	67
Pepper-mint	98
Pepper Water	121
Periwinkle	183
greater	37
Pestilent Wort	122
Petty Whin	9
Pig-nut or Hog-nut	20
Pilewort	34
Pimpernel, male	8
Pin Pealwort	7
Pink Deptford	29
Meadow	90
Pipperidge Bush	15
Plaintain, Buck's horn	124
great	125
hoary	125
ribwort	125
Water, great	124
star-headed	43

	Page
Polypody	127
branched	53
cristed	53
female	52
Pond-weed, broad leaved	131
fine or fennel leaved	130
horned	67
long leaved, great, with pel-	
lucid leaves	129
perfoliated	131
Poplar Tree, black	128
white	128
trembling	128
Poppy, bastard	116
Corn	115
long, rough-headed	116
red	115
round, smooth-headed	115
Prick-madam	152
Prickwood, Cornel-tree	39
Spindle-tree	49
Primrose, common	132
Privet or Prim	86
Pudding-grafs	134
Purflane, Water	129
Pursley-Piert	119

Q.

	Page
Quick	100
Quicken Tree	162

R.

Radish, Horse	141
Water	136
Ragwort, broad leaved marsh	72
common	72
hoary perennial with Groundsel leaves	72
Rattle, yellow	117
Reed Mace	175
Rest-harrow	9
Rib-wort Plantain	125
Roan Tree	162
Rocket, base	142
or Winter Cresses	14
Rosa Solis, or Sun-dew	143
Rose-bay, Willow herb	94
Rose, dogs red flowered	144
white flowered	143
Rue, meadow	166
whitlow grass	150
Rupture-wort, the least	136
Rush, flowering, or Water Gladiole	20

Sage,

S.

Page

Sage, Wood	151
Saint-foin	109
Saint John's wort	68
hairy	68
least trailing	69
upright	70
Saint Peter's wort	69
Sallow, common round leaved	148
Sanicle	149
Satyrion Royal, male handed	111
female handed	111
Sauce alone	64
Saw-wort	155
Saxifrage, common golden	149
Meadow	155
Scabious common Field double flowered	150
lesser Field	150
Scorpion-grass	102
small yellow flowered	102
Water	102
Self-heal	132
Sengreen	153
Septfoil	169
Service Tree, common wild	99
Setterwort	64
Shepherd's Needle	151
Purse	20
Rod	45
	Silver

Page

Silver Weed	119
Simson	154
Sloe Tree	133
Smallage	10
Snapdragon, the greater	11
the least	10
Sneezewort	134
Soapwort	90
Solomon's Seal	191
Sorb Tree	99
Sorrel, common	80
Sheeps	80
Wood	114
Sow-bane	17
Sow-thistle, common	161
prickly	161
with less jagged leaves	161
smooth with less jagged leaves	160
Tree	161
Spear-thistle	24
Spearwort-leffer	139
Speedwell, or Chickweed	181
ivy leaved	181
little smooth	182
male	182
Water, narrow leaved	179
Spatling Poppy	89
Spindle Tree	49
Spleenwort, or Miltwaft	12
rough	88
2	Spurge,

	Page
Spurge, broad leaved	169
dwarf, small annual	168
Laurel	84
petty	169
rough fruited	168
Sun, or Wart wort	168
Wood	167
Spurrey, Corn	7
purple	163
Star-headed Water Chickweed	163
of the earth	124
Thistle	26
Star-wort, autumnal	164
vernal	163
Stitchwort, greater	27
least	7
lesser	28
a variety	28
Stonecrop, yellow	152
Wall Pepper	153
Stone Parsley, bastard	158
Strangle-Tare	183
Strawberry	54
barren	55
trefoil	171
Succory, wild	35
Sun-dew, round leaved	143
Sweet-briar	144
Swine's Cresses	105

T.

	Page
Tansey, common	165
wild	119
Tare, everlasting	83
hairy	40
small wild	40
smooth	41
strangle	183
tine	40
with small pods	41
Teasel, small wild	45
wild	45
Tentwort	145
Thistle, common Way, or creeping	26
wild Carline	27
cotton	26
marsh	25
Milk, or Lady's	25
Spear	24
Star	26
upon Thistle	24
welled, with small flowers	25
Thorn-Apple	159
Thorn black	133
white	100
common purging	143
Three faces under a hood	185
Throatwort, little or lesser	23
Thyme, Mother of	154
	Tine

	Page
Tine Tare	40
with small pods	40
Toad Flax, common yellow	87
ivy leaved	87
Tormentil	169
Tower Mustard, great	174
Traveller's Joy	37
Trefoil, Bird's-foot	88
greater	88
common purple	174
Hare's-foot	171
Heart	171
Hop	173
lessor, or decumbent	172
knotted	173
long leaved purple	174
Marsh	97
Melilot	172
small	172
Strawberry	171
white	170
flowered, meadow	173
Trichomanes	170
Triple Ladies-Traces	112
Tun-hoof	22
Tutsan	68
hairy	68
Tway-blade, common	190
Two faces under a hood	183

V.

	Page
Valerian, great wild	176
jagged leaved	76
small, wild, or Marsh	176
another sort	177
Venus-comb	151
looking-glass, lesser	22
Vervain	178
mallow	4
Vetch Bush	182
tufted	40
wild	183
Vetchling, common yellow bastard	83
Medick	109
Violet, Dog's or wild	184
purple sweet	184
Throat wort leaved	184
Viper's Bugloss	46

W.

Wake Robin	11
Wall-flower	85
pepper	153
rue	145
wort	149
Wart-wort	168

Water-Betony	152
Caltrops greater	130
leffer	130
Crow-foot fennel leaved	137
fine leaved	138
finest leaved	137
Ivy leaved	138
round leaved	140
various leaved	137
Creffes	157
Dock, great	79
Dropwort	108
Hemlock	109
leffer	108
Elder	110
Figwort	152
Gladiole	20
Hemlock	122
Hemp agrimony, trifid	179
with an undivided	
leaf	179
Horehound	91
Lily, white	106
yellow	106
leffer, with a fringed	
flower	107
Mill-foil, feathered or spiked	129
verticillated	119
Mint	97
a variety	157
with whorled coronets	98
Water	

	Page
Water Parsnep, creeping	159
great	158
least	158
Pepper	121
Plantain, great	124
Star-headed	43
Purslane	129
Radish	136
Ragwort	72
Speedwell, middle long-leaved	180
narrow leaved	179
Wort	163
Way-bread	125
Weed, yellow	89
Whin	58
petty	58
White-rot	67
White-thorn	100
Whitlow-grafs, common	116
rue-leaved	150
Wilding	96
Williams, wild	90
Willow, common white	147
creeping dwarf	148
round eared shining, or Almond	
leaved	147
sweet	190
Willow-herb	146
great flowered	93
greater smooth leaved	92
hooded	29
least smooth leaved coddled	93
Willow-	

	Page
Willow-herb, leffer hairy coddred with small flowers	94
middle smooth leaved	92
rose bay	94
yellow	92
Winter-creffes, or rocket	49
Woad, wild	89
Wood Anemone	8
Betony	15
Wood-bine	23
Woodroof	12
Wood Pea, or Heath Pea	113
Sorrel	114
Wood-waxen	59
Sage	151
Spurge	167
Wormwood, common	1
Wych-Elm	187
Hazel	188

Y.

Yarrow common	100
Yellow-weed, common	89
Yew Tree	166


E R R A T A.

In printing this CATALOGUE, some Errors have escaped the Press. In a few Instances, the English Name of the Plant is printed in Italick Characters: some Errors occur in the Pointing; and the Greek Letter, by which a different Species is marked in *Hudson's FLORA ANGLICA*, is not always in its proper Place. These which follow, with any others that may occur, are left to the candid Reader, to excuse and correct.

page	line		
3	15	for Tetrandia	read Tetrandia
		and in other Places,	where this <i>Erratum</i> is found.
4	4	for Monodelphia	read Monadelphina
10	21	Orentium	Orontium
11	27	Point	Pint
24	13	Polygamiara	Polygamia
31	13	dichotimo	dichotomo
32	11	temulum	tremulum
34	22	longissimis	longissimis
37	2	Lutesiana	Lutetiana
—	23	foilis primatis	foliis pinnatis
39	3	peniculato	paniculato
43	4	hexaginis	hexagynis
—	8	the same	
50	9	Euparasia	Euphrasia
53	3	sub pinnatis	sub bipinnatis
—	20	bipennata	bipinnata
56	11	pericrapiis	pericarpis
60	19	Leighton	Layton
64	19	Echoides	Echioides
—	21	Echinoides	Echioides
—	—	involuero	involucro
76	1	aguina	agnina
79	21	Polygyamia	Polygamia
83	19	lutens	luteus
84	14	and	and
86	3	Diandrya	Diandria
89	25	Popy	Poppy
94	5	sub simplici	subsimplici
99	19	terminalis	torminalis
102	21	Moufe ear	Water
104	1	Syvestris	Sylvestris
112	10	eight	ten

E R R A T A.

page	line		
117	23	<i>for</i> Coxcomb	<i>read</i> Cock's comb
—	22	angiospetmia	angiospermia
118	6	the same	
128	8	Kenis	Kenis
122	—	Wert	Wort
123	14	Saxifraga	Saxifrage
124	5	Monadaiphia	Monadelphia
—	15	on	in
125	10	Plantaco	Plantago
131	18	scapho	scapo
137	25	various water leaved	water various leaved
		crowfoot	crowfoot
139	18	nemerosus	nemorosus
141	26	<i>after</i> Water Radish,	<i>add,</i> with jagged leaves
150	—	<i>for</i> Monognyia	<i>read</i> Monogynia
152	—	Scrophulalia	Scrophularia
159	2	Axilaribns	Axillaribus
164	3	Callatriche	Callitriche
172	4	persistentibus	persistentibus
180	22	recemosis	racemosis



I N D E X

O F T H E

L A T I N N A M E S,

A S G I V E N

By L I N N Æ U S.

A.

	Page
Acer campestre	2
Pseudo platanus	2
Achillea millefolium	100
Ptarmica	134
Acorus Calamus	2
Aegopodium Podagraria	9
Aethusa Cynapium	36
Agrimonia Eupatoria	3
Agrostemma Githago	90
Ajuga reptans	19
Alisma Damasonium	43
Plantago	124
Allium vineale	4
Amarantus Blitum	18
G g	Anagallis

	Page
Anagallis arvensis	8
Anemone nemorosa	8
Anethum Foeniculum	54
Angelica sylvestris	9
Anthemis Cotula	32
nobilis	33
Antirrhinum Cymbalaria	87
Elatine	86
Linaria	87
majus	11
orentium	10
Aphanes arvensis	119
Apium grave olens	10
Arctium Lappa	82, 83
Arenaria rubra	163
serpyllifolia	6
trinervia	7
Artemisia Absinthium	1
vulgaris	11
Arum maculatum	11
Asperugo procumbens	19
Asperula odorata	12
Asplenium Adiantum nigrum	3
Ceterach	12
Ruta muraria	145
Scolopendrium	122
Trichomanes	170
Atriplex hastata	13
patula	12
Atropa Belladonna	14
B.	
Ballote nigra	14
Bellis perennis	15
Berberis vulgaris	15
	Beotnica

Page

Betonica officinalis	15
Betula alba	16
Alnus	4
Bidens tripartita	179
Borago officinalis	18
Brassica Napus	104
Bryonia alba	18
Bunium Bulbocastanum	29
Butomus umbellatus	20
Buxus sempervirens	21

C.

Callatriche autumnalis	164
verna	163
Caltha palustris	127
Campanula glomerata	23
hybrida	22
rotundifolia	23
Cardamine pratense	24
Carduus acanthoides	25
crispus	24
lanceolatus	24
Marianus	25
palustris	25
Carlina vulgaris	27
Carpinus Betulus	114
Caucalis anthriscus	30
arvensis	30
Centaurea Calcitrapa	26
Cyanus	41
Jacea	71
Scabiosa	71

	Page
Cerastium aquaticum	6
semidecandrium	32
viscosum	5
vulgatum	5
Ceratophyllum demersum	67
Chaerophyllum sylvestre	36
tremulum	32
Cheiranthus Cheiri	85
Chelidonium majus	115
Chenopodium album	16
Bonus Henricus	17
hybridum	35
murale	17
polyspermum	34
rubrum	17
serotimum	16
urbicum	34
Chrysanthemum Leucanthemum	85
segetum	35
Chrysosplenium oppositifolium	149
Cichoreum Intybus	35
Circoea Lutetiana	37
Clematis Vitalba	37
Clinopodium vulgare	38
Cochlearia armoracia	141
Coronopus	105
Conium maculatum	36
Convallaria majalis	86
multiflora	191
Convolvulus arvensis	38
sepium	38
Cornus sanguinea	39
Corylus Avellana	40
Cratægus	

Page

Cratægus Oxycantha	100
terminalis	99
Crepis tectorum	65
Cucubulus Behen	89
Cuscuta Europaea	41
Cynoglossum officinale	42

D.

Daphne Laureola	84
Datura Stramonium	159
Daucus Carota	43
Delphinium Consolida	44
Dianthus Armeria	29
Digitalis purpurea	45
Dipsacus pilosus	45
sylvestris	45
Draba verna	116
Drosera rotundifolia	143

E.

Echium vulgare	46
Epilobium angustifolium	94
hirsutum	94
montanum	92
palustre	93
ramosum	93
tetragonum	92
Equisetum arvense	46
palustre	47
sylvaticum	47
Erica cinerea	48
Tetralix	47
vulgaris	48

Ervum

	Page
Ervum hirsutum	40
tetraspermum	41
Erysimum Alliaria	64
Barbarea	14, 49
officinalis	48
Euonymus Europaeus	49
Eupatorium Cannabinum	50
Euphorbia Amygdaloides	167
exigua	168
Helioscopia	168
Peplus	169
platyphyllos	169
verrucosa	168
Euphrasia officinalis	50
odontites	50

F.

Fagus Castanea	29
sylvatica	51
Ficaria verna	34
Filago Germanica	62
montana	61
Fragaria sterilis	55
vesca	54
Fraxinus excelsior	55
Fumaria capreolata	56
officinalis	56

G.

Galeopsis Ladanum	156
Tetrahit	77, 78
Galeobdolon	78
Galum Aparine	10
	Gallium

Gallium verum	57
Mollugo	101
montanum	101
palustre	101
Genista Anglica	58
tinctoria	59
Gentiana Centaurium	31
Geranium cicutarium	59
dissectum	60
molle	60
pusillum	60
Robertianum	61
Geum urbanum	27
Glechoma arvensis	156
hederacea	22
Gnaphalium uliginosum	61

H.

Hedera Helix	63
Hedysarum Onobrychis	109
Helleborus foetidus	64
Heracleum Sphondylium	163
Hieracium Pilosella	123
fabaudum	65
Humulus Lupulus	89
Hyacinthus non-scriptus	67
Hydrocotyle vulgaris	67
Hyoscyamus niger	68
Hypericum Androsaemum	69
hirsutum	68
humifusum	69
perforatum	68
pulchrum	70
quadrangulum	69

Hypo-

	Page
Hypochaeris radicata	65
I.	
Ilex Aquifolium	3
Inula dysenterica	39
Helenium	63
Pulicaria	39
Juniperus communis	74
Iris foetidissima	73
Pseudacorus	73
L.	
Lactuca virofa	76
Lamium album	77
Amplexicaule	78
rubrum	79
Lampfana communis	79
Lathyrus Nissolia	30
pratensis	83
Lemna minor	84
polyrrhiza	85
trifulca	84
Leontodon autumnale	66
hispidum	44
Taraxacum	44
Ligustrum vulgare	86
Linum catharticum	87
Radiola	136
Lithospermum arvense	19
Lithrum Salicaria	146
Lonicera Periclymenum	23
Lotus corniculata	88

	Page
Lychnis dioica	91
Flos cuculi	90
Lycopus Europæus	91
Lyfimachia Nummularia	105
tenella	106
vulgaris	92
M.	
MalvaAlcea	4
rotundifolia	95
fylvestris	95
Matricaria Chamomilla	33
inodora	32
Parthenium	96
Medicago Arabica	171
lupulina	172
Melampyrum fylvaticum	96
Mentha aquatica	97
arvensis	98
piperita	98
Pulegium	134
Menyanthes Nymphoides	107
trifolia	97
Mercurialis annua	99
perennis	99
Montia fontana	6
Myrica Gale	190
Myosotis scorpioides	142
Myosurus minimus	103
Myriophyllum spicatum	129
verticillatum	119
N.	
Narcissus Pseudo-Narcissus	104
Nepeta Cataria	105
H h	Nymphæa

	Page
Nymphæa alba	106
lutea	106
O.	
Oenanthe crocata	109
fistulosa	108
Ononis spinosa	9
Onopordum Acanthium	26
Ophioglossum vulgatum	109
Ophrys ovata	190
spiralis	112
Orchis bifolia	110
latifolia	111
macula	110
maculata	111
morio	111
Origanum vulgare	112
Ornithopus-perpusillus	112
Orobanche major	113
Orobis tuberosus	113
Osmunda regalis	53
spicant	88
Oxalis Acetosella	114
P.	
Papaver Argemone	116
Rhoeas	115
Parietaria officinalis	116
Pastinaca sativa	117
Pedicularis palustris	117
sylvatica	118
Peplis Portulaca	129
Pteris aquilina	52
Phellandrium aquaticum	122
Picris	

	Page
Picris Echinoides	64
Pimpinella saxifraga	123
Pinus sylvestris	124
Plantago Coronopus	124
lanceolata	125
major	125
media	125
Polygala vulgaris	126
Polygonum amphibium	121
aviculare	126, 127
Convolvulus	51
Hydropiper	121
pensylvanicum	121
Perficaria	120
Polypodium aculeatum	52
cristatum	53
Dryopteris	53
Filix mas	54
vulgare	127
Populus alba	128
nigra	128
tremula	128
Potentilla argentea	118
Argentina	119
reptans	118
Potamogeton crispum	130
lucens	129
pectinatum	130
persoliatum	131
natans	131
ferratum	130
Primula veris	131, 132
vulgaris	132
Prunella vulgaris	132

	Page
Prunus infititia	133
Padus	31
spinosa	133
Pyrus communis	134
Malus	96
Q.	
Quercus Robur	135
R.	
Ranunculus acris	140
aquatilis	137, 138
arvensis	139
auricomus	139
bulbosus	138
Flammula	139
hederaceus	138
repens	140
sceleratus	140
Raphanus Rraphanistrum	141, 142
Reseda lutea	142
Luteola	89
Rhamnus catharticus	143
Frangula	55
Rhinanthus Crista galli	117
Rosa arvensis	143
canina	144
Eglanteria	144
Rubus caesius	145
fruticosus	144
Rumex Acetosa	80
Acetosella	80
acutus	80
Britannica	79
crispus	81
obtusifolium	82
	Ruscus

Ruscus aculeatus	145
Sagina erecta	7
procumbens	7
Sagittaria Sagittifolia	146
Salix alba	147
amygdalina	147
caprea	148
fusca	148
vimenalis	147
Salvia Verbena	66
Sambucus Ebulus	149
nigra	148
Sanicula Europaea	149
Saponaria officinalis	90
Saxifraga trydactylites	150
Scabiosa arvensis	150
Columbaria	150
succisa	151
Scandix Anthriscus	103
Pecten Veneris	151
Scleranthus annuus	75
Scrophularia aquatica	152
nodosa	152
Scutellaria galericulata	29
Sedum acre	153
reflexum	152
Telephium	83
Sempervivum tectorum	153
Senecio aquaticus	72
erucifolius	72
Jacobaea	72
sylvaticus	154
viscosus	153
vulgaris	154
	Serra-

	Page
Serratula arvensis	26
tinctoria	155
Seseli Caruifolia	155
Sinapis alba	156
arvensis	142
nigra	157
Sison Amomum	158
inundatum	158
Sisymbrium amphibium	136, 141
Nasturtium	157
Sophia	49
Sium latifolium	158
nodiflorum	159
Solanum Dulcamara	159
nigrum	160
Solidago Virga aurea	185
Sonchus arvensis	161
oleracius	160, 161
Sorbus aucuparia	162
Sparganium erectum	162
Spartium scoparium	58
Spergula arvensis	7
Spiraea Filipendula	52
Ulmaria	187
Stachys palustris	155
sylvatica	57
Stellaria graminea	5
Holothea	27
Symphytum officinale	164

T.

Tamus communis	165
Tanacetum vulgare	165
Taxus baccata	166
Teucrium Scorodonia	151
Thalictrum	

	Page
Thalictrum flavum	166
Thlaspi Bursa Pastoris	29
campestre	166
Thymus Serpyllum	154
Tilia Europaea	167
Tordylium nodosum	31
Tormentilla erecta	169
Tragopogon pratense	170
Trifolium agrarium	173
arvense	171
filiforme	172
fragiferum	171
medium	174
Melilotus officinalis	97
pratense	174
procumbens	172
repens	170, 173
striatum	173
Tussilago Petasites	122
Farfara	175
Turritis glabra	174
Typha latifolia	175

V.

Valeriana dioica	176, 177
officinalis	176
Locusta	76, 177
Verbascum nigrum	178
Thapsus	178
Verbascina cernua	179
Verbena officinalis	178
Veronica agrestis	181
Anagallis	180
arvensis	181
3	Veronica

	Page
Veronica Becabunga	180
Camaedrys	180
Hederifolia	181
officinalis	182
serpyllifolia	182
scutellata	179
Viburnum Opulus	110
Vicia Cracca	40
Lathyroides	183
sepium	183
Vinca major	37
minor	183
Viola canina	184
hirta	184
odorata	184
tricolor	185
Viscum album	186

U.

Ulex Europaeus	58, 59
Ulmus campestris	187, 188
glabra	188
Urtica dioica	189
urens	188







